SMALLEST ANNUAL WEEDS prepared by: Zharkynbekkizy A Mambetaliyeva M Adopted by:

PLAN

- 1. The concept of the weeds
- 2. The dangers of
- 3. Classification of weeds and their common types
- 4. Smallest annual weeds
- 5. Ephemeral
- 6. Winter weeds
- 7. Biennial weeds
- 8. Perennial weeds
- 9. Umbilical vascular weeds
- 10. The root system of fringe weeds

The concept of the weeds

Weed plants growing in the fields of culture that is detrimental to the quality of their products to reduce the amount of plants. Cultural crop rotation and other cultural crops is encountered, it is called the polluting plants. Crop growing faster than weeds, spend a lot more moisture and nutrients..



The dangers of

 Shade crop of weeds growing on the top floor, which slows down the process of photosynthesis in the leaves, accompanied by the deterioration of agricultural machines, which will cause the class. agricultural machines in their class can cause deterioration of the work.







Classification of weeds and their common types

 Nearly 300 species of weeds. Patrolling the classification based on several criteria: power plants method, duration of survival, reproduction method. The method of power-minded, semi-minded and not-minded weeds off.

Smallest annual weeds

 Life expectancy are divided into 2 groups: one year, a lot of the year. One-year survival of weeds process lasts from 1 to 2 years. Their increasing appearance of new growths and early summer, depending on the length of the stages of development of these ephemeral, late summer, winter and the year is divided into two.





Ephemeral

 Ephemeral stages of growth and development in a very short (1.5-2 months) plants. They can be several generations per season. In this group, the most harm to black oats.





Winter weeds

For the development they need to lower the temperature of the winter season. Despite the terms of new growths, the plants will spend the winter, and fruit in the summer of next year. With winter cultural crops grow and mature at the same time the family. Representatives: rye, barley field.

Biennial weeds

 Plants to complete within two years of their life. First formed in the root system and leaves the stork. Representatives of this group meñdwana black, white and yellow camel alfalfa, etc.



Perennial weeds

They grow a few years. Only members of the surface of the soil in the winter dies. Winter roots will continue to grow next year. They can grow and go through the veins through the family. Representatives of the main field ivy, creeping repens, fiber spurge, etc.



Thank you for attention!!!