

# Cognitive Behavioral Treatment of Bipolar Disorder

The original version of these slides was provided by

Michael W. Otto, Ph.D.

with support from NIMH Excellence in Training Award at  
the Center for Anxiety and Related Disorders at Boston  
University

(R25 MH08478)

# Use of this Slide Set

- Presentation information is listed in the notes section below the slide (in PowerPoint normal viewing mode).
- References are also provided in note sections for select subsequent slides

# Diagnostic Considerations

- Manic Episode
  - 1 week high, euphoric, or irritable mood plus 3 (4) of the following:
    - ◆ exaggerated feelings of importance
    - ◆ little need for sleep
    - ◆ racing thoughts
    - ◆ pressured speech
    - ◆ distractibility
    - ◆ increased goal directed behavior (agitation)
    - ◆ reckless behavior
- Hypomanic Episode
  - 4 days of high, euphoric, or irritable mood plus 3 (4) symptoms (no impairment, psychotic features, need for hosp.)


# Diagnostic Considerations

- Bipolar I
  - At least one manic or mixed episode
  - May or may not have depressive episode, but most do (71% of sample)
  - 3.5 more likely to have depressive symptoms than manic/hypomanic (Judd et al., 2002)
- Bipolar II
  - At least one hypomanic episode and one or more depressive episodes
  - 38 times more likely to have depressive symptoms than hypomania (Judd et al., 2003)
- Bipolar I vs II status is only inconsistently predictive of shorter term outcomes (cf., Judd et al., 2003; Miklowitz et al., 2007; Otto et al., 2006).

# Characteristics of Patients With Bipolar Disorder

- Prevalence
  - 1-2% of the population
- Age of Onset
  - Late teens to early 20s (earlier age of onset is associated with a worse course; Perlis et al. 2006).
- Sex Ratio
  - Equal, but more rapid cycling among women
- Comorbidity
  - Anxiety, Substance Use, ADHD
- Course
  - 75% relapse 4-5 years, half in 1 year (the proportion of days ill predicts episode frequency the next year; Perlis et al., 2004)

# Comorbidity in Bipolar Disorder (assessed in 1000 patients enrolled in STEP-BD)



Kogan et al., 2004

# Diagnostic Issues – Major Depression

- Depression:
  - Youth hospitalized for severe depression (young and severe) – 41% experienced manic/hypomanic episode over next 15 years (Goldberg et al., 2001)
- Depression + Substance Use Disorder
- Depression + Borderline Disorder
- Depression + psychosis (schizoaffective disorder)

# Presentation with Psychosis

- Is it mania?
- Schizophrenia?
- Substance Induced?
- Schizoaffective?
  
- History and family help



# An Abundance of Distress and Disability

- Family, job, personal
- Post-episode studies
  - 6 months after: 30% unable to work; only 21% worked at their expected level (Dion et al., 1988)
  - 1.7 years after hospitalization: 42% had steady work performance (Harrow et al., 1990)
- Relatively high rates of suicide in bipolar disorder (predicted prospectively by days depressed and previous attempts; Marangell et al., 2006)

# Psychosocial Treatment

# Topics

- What is the evidence for the efficacy of psychotherapy for bipolar disorder?
- What are the targets of treatment?
- What are the elements of treatment?

# Psychosocial Treatment for Bipolar Disorder

## Initial Encouragement:

- Psychosocial Predictors of Bipolar Course
- Incomplete Efficacy of Mood Stabilizers
- Practice Characteristics
  - Majority of bipolar patients are engaged in some sort of psychosocial care

## Direct Evidence

- Promising outcomes from well-controlled trials

# Role of Psychosocial Factors in Bipolar Disorder

- Psychosocial stressors impact the course of bipolar disorder:
  - Family stress (expressed emotion)<sup>1</sup>
  - Negative life events<sup>2</sup>
  - Cognitive style<sup>3</sup>
  - Sleep disruptions<sup>4</sup>
  - Anxiety comorbidity<sup>5</sup>

<sup>1</sup> Miklowitz et al. (1988)

<sup>2</sup> Johnson & Miller, (1997); Ellicott et al. (1990)

<sup>3</sup> Reilly-Harrington et al., 1999

<sup>4</sup> Malkoff-Schwartz et al. (1998)

<sup>5</sup> Simon et al. (2004); Otto et al. (2006)

# Pharmacotherapy for Bipolar Disorder

- Advances in the field, but among patients taking medications:
  - Half relapse first year
  - Three-quarters relapse over several years
  - Continued role impairment between episodes
  - Poor medication adherence

(Gitlin et al., 1995; Keck et al., 1998; O'Connell et al., 1991; Tohen et al., 1990)

# Focused Psychosocial Treatments for Bipolar Disorder

- The product of diverse theoretical orientations, but with a high degree of similarity in strategies.
- In particular, randomized trials have shown support for
  - Cognitive Behavioral Therapy (CBT)
  - Interpersonal and Social Rhythm Therapy (IPSRT)
  - Family-Focused Treatment (FFT)

# Common Treatment Elements Among CBT, IPSRT, FFT

- Psychoeducation providing a model of the disorder and risk and protective factors (e.g., the role of sleep and lifestyle regularity).
- Communication and problem-solving training aimed at reducing familial, relationship, or external stress.
- Review of strategies for the early detection and intervention with mood episodes (including increased support, pharmacotherapy, more-frequent monitoring).



# Some of the Influential, Psychosocial Clinical Trials

- Medication adherence<sup>1</sup>
- Detection of prodromal episodes, early intervention<sup>2</sup>
- Individual CBT for Relapse Prevention<sup>3</sup>
- Individual IPSRT for Relapse Prevention<sup>4</sup>
- Family Interventions for Relapse Prevention<sup>5</sup>
- Group Psychoeducation for Relapse Prevention<sup>6</sup>
- Individual CBT for Episode Treatment<sup>7</sup>
- Intensive CBT, IPSRT, or FFT for Bipolar Depression<sup>8</sup>

1. Cochran (1983)

2. Perry et al. (1999)

3. Lam et al. (2000), Lam et al. (2003); Scott et al. (2001)

4. Frank et al. (1997); Frank et al. (1999)

5. Miklowitz et al. (2003); Rea et al. (2003); Simoneau et al. (1999); also Clarkin et al. (1998)

6. Colom et al. (2003)

7. Scott et al. (2006)

8. Miklowitz et al. 2007

# Cognitive-Behavioral Therapy (CBT) for Medication Adherence (Cochran, 1984)

## Relapse Prevention


- 6 sessions of adjunctive CBT vs standard clinical care<sup>4</sup>
- At end point and at 6-month follow-up, CBT patients had
  - Greater medication adherence
  - Lower hospitalization rates

Cochran S. *J Consult Clin Psychol.* 1984;52:873-878.

# Lam et al. - An Early CBT Success


- 103 bipolar patients randomized to CBT or TAU
- 12-18 sessions individual CBT
  - Information
  - Monitoring of mood & cognitions (early intervention)
  - Management of sleep and routine
  - Attention to “making up for lost time”
- 8 dropout in each condition

# Medication Adherence




# Clinical Outcome (days ill over 1 year)

Mean Days in Status



# Survival Analysis

(N = 103)



# Family-Focused Treatment

## Elements


- Psychoeducation about bipolar disorder
- Communication-enhancement training
- Problem-solving training<sup>1</sup>

## Outcome

- Adjunctive FFT appears to effect<sup>1</sup>
  - Depressive symptoms
  - Manic symptoms
  - Rehospitalization times

# 1-Year Survival Rates Among Bipolar Patients in Family-Focused Treatment versus Case Management

Miklowitz DJ, et al. *Arch Gen Psychiatry*. 1988;45:225-231.



Wilcoxon Test,  $\chi^2 (1) = 4.4, p = .035$



# Six Objectives of FFT

Help the patient and her or his relatives to:

- Understand the nature of bipolar disorder and cyclic mood disturbances.
- Accept the concept of vulnerability to future episodes
- Accept a crucial role for mood-stabilizing medication for symptom control
- Distinguishing between personality and bipolar disorder
- Recognize and develop coping skills for managing the stressful life events that trigger recurrences of bipolar disorder
- Reestablishing role and interpersonal functioning after a mood episode

# Interpersonal and Social Rhythm Therapy

- Educate patient about bipolar disorder
- Identify current interpersonal problem areas (e.g., grief, disputes, role transitions, interpersonal deficits)
- Initiate social rhythm metric

Frank et al. *Biological Psychiatry* 1997 1165-1173

# Group Psychoeducation vs. Standard Care


- 21 Weeks of Randomized Treatment, 2-year follow-up
- 120 outpatients in remission for 6 months
- Standard Care
  - Treatment algorithms
  - Monthly sessions
  - Serum levels of medications assessed
- Group Treatment 21 90-minute sessions
- Outcome
  - Recurrences at endpoint: 38% in group vs. 60% in SC
  - Recurrences at 2 years: 67% in group vs. 92% in SC

# Psychoeducation?


- Psychoeducation
  - What is bipolar illness
  - Symptoms
  - Treatments
  - Serum levels
- Early detection of episodes
- Risk reduction - substance use
- Lifestyle regularity
- Stress management
- Problem solving

# CBT, IPSRT, FFT vs. Collab Care for Bipolar Depression

Miklowitz et al., 2007, Archives Gen Psychiatry



# No Significant Differences Among the Intensives: CBT, IPSRT, FFT



# Given this Evidence...

## ...What are Some Targets for Psychotherapy?

- Medication adherence
- Early detection and intervention
- Stress and lifestyle management
- Treatment of bipolar depression
- Treatment of comorbid conditions

# Medication Non-Adherence in Mood Disorder

- 98 patients taking mood stabilizers (80% bipolar)
- 50% non-adherence rate last year
- 30% non-adherence last month ( $\leq 70\%$  adherent)
- Predictors of non-adherence:
  - denial of severity of illness
  - previous non-adherence
  - greater illness duration

(Scott & Pope, 2002, J Clin Psychiatry, 63:384-390)



# Relapse Prevention

- Patient as cotherapist
- Treatment contract
- Training in early detection
- Use of treatment team

# Individualized Treatment Contract

- Why contract?
- Formulate a plan for the future
- How I know I am depressed
  - Plan during depression
- I am manic when...
  - Plan during mania (include who initiates the plan)
- Other modules
  - Substance abuse, Bulimia, Gambling, Budget, etc

# Mood Charting

- Enables early and accurate identification of changes in mood
- Allows for early intervention prior to severe episodes
- Tracks medication doses and adherence to psychological treatment
- Tracks hours slept and sleep/wake times
- Notes daily psychosocial stressors that may serve as triggers for relapse

# Strategies for Hypomania

- Explore medical solutions (e.g., dosage or medication changes)
- Counteract impulsivity
  - Give car keys or credit card to someone to hold
  - “Make rules” about staying out late or giving out phone number
  - Avoid alcohol and substance use
- Minimize stimulation
  - Avoid confrontational situations

# Cognitive-Behavioral Therapy for Bipolar Depression/Relapse Prevention

## Structure of Sessions

- Review of symptoms, progress, and problems
- Construction of the agenda
- Discussion, problem solving, rehearsal
- Consolidation of new information/strategies
- Assignment of home practice
- Troubleshooting of homework (including signposts of adaptive change)

# Cognitive Restructuring and Skill Acquisition

## Restructuring

- Education (role and nature of thoughts)
- Self-monitoring of thoughts
- Identification of errors
- Substitution of useful thoughts
- Core beliefs and strategies

## Skill acquisition

- Assertiveness
- Communication skills
- Problem solving

# Cognitive Restructuring

---

- ◆ Examine the evidence for the thought
  - ◆ Generate alternative explanations
  - ◆ De-catastrophize
  - ◆ Debunk “shoulds”
  - ◆ Find the logical error
  - ◆ Test out its helpfulness
-

# Questions Used to Formulate Rational Response

---

- ◆ What is the evidence that the automatic thought is true? Not true?
  - ◆ Is there an alternative explanation?
  - ◆ What is the worst that could happen? Would I live through it?
  - ◆ What's the best that could happen?
  - ◆ What's the most realistic outcome?
-



# Questions Used to Formulate Rational Response (Cont'd)

---

- ◆ What is the effect of my believing the automatic thought?
  - ◆ What is the cognitive error?
  - ◆ If a friend was in this situation and had this thought, what would I tell him/her?
-

# Respecting Hot Emotions

- Interventions are in relation to, not in spite of, the patient's current mood.
- Train emotional regulation skills
- Gain access to mood-state dependent cognitions

# Activity Assignments: Bipolar Disorder

- Management of sleep
- Management of over/under activity
- Management of destructive activities (substance use)
- Resetting goals given limitations due to the disorder

# Activity Assignments - 1

- Independent Intervention or used in conjunction with cognitive restructuring
- Help ensure that therapy is not over-focused on thinking rather than doing
- Often requires a problem-solving analysis to understand patterns of over- and under-activity relative to the patient's values

## Activity Assignments - 2

- Monitor current Activities
- For change:
  - Start small (where the patient is)
  - Be specific
  - Rehearse elements in session
  - Define outcome objectively
  - Troubleshoot problems and signposts
  - Review cognitions (expectations, concerns)

## Activity Assignments - 3

- Review performance relative to objective criteria (and the degree of mood disturbance)
- Assess the patient's cognitive and emotional response to the assignment
- Discuss further applications

# Well-Being Therapy Phase

- In this phase, therapeutic effort and monitoring is devoted to increasing periods of well being rather than reducing pathology.
- It provides a way to consolidate gains around positive outcomes
- An excellent strategy for fading out treatment

# End of Treatment

- Patient has skills to act as his or her own therapist
- Patient focuses on well-being
- Therapist contact fades



# Cognitive-Behavioral Therapy for Comorbid Disorders

- Anxiety disorders
- Substance use disorders
- Eating disorders

# New Directions in CBT for Bipolar Disorder

## Promoting Emotional Tolerance

- Getting better with the rollercoaster of emotions
- Learn to apply emotional acceptance plus problem solving in the context of strong emotions (anxiety, sadness, euphoria)
- Initial evidence for mindfulness training in bipolar disorder – improvements in mood and cognitive symptoms
  - ◆ (Deckersbach et al., 2012, CNS Neurosci Ther).