



УНИВЕРСИТЕТ
ЛОБАЧЕВСКОГО

Lecture 7


REGULATION OF ENZYME ACTIVITY AND METABOLIC PATHWAYS. MEDICAL IMPLEMENTATION OF ENZYMES

Dr. Elena O. Polovinkina
2020


Ways of regulation of enzymatic activity




УНИВЕРСИТЕТ
ЛОБАЧЕВСКОГО
НИЖЕГОРОДСКИЙ ГОСУДАРСТВЕННЫЙ УНИВЕРСИТЕТ




Enzymes phosphorylation/ dephosphorylation




The scheme of activation of pepsinogen – partial proteolysis



Dissociation of protein kinase A - Protein-protein interactions




Allosteric regulation



Regulation of the metabolic pathways




The key enzymes in the metabolic pathways usually catalyze:

- ▶ irreversible (\rightarrow) or partially reversible reactions;
- ▶ the slowest reactions;
- ▶ reactions at the beginning of the metabolic pathways or in branching places of metabolic pathways.



Scheme of regulation of enzyme activity by feedback inhibition

Regulation of the metabolic pathways



Clinical applications of enzymes

- ▶ enzyme diagnostics;
enzymopathology:
 - accumulation of substrate reaction. For example, phenylalanine in phenylketonuria, free bilirubin in physiological jaundice of the newborn, some fats in diseases of lysosomal accumulation (lipidosis);
 - product deficiency. For example, melanin in albinism, catecholamines in parkinsonism;
 - both features simultaneously, as in glycogenosis, accompanied by hypoglycemia with an excess of glycogen in the liver;
- ▶ enzyme therapy;
- ▶ use of enzymes in medical technology and industry (Enzyme immunoassay: *enzyme-antigen-antibody*);
- ▶ use of enzyme inhibitors:
 - natural and artificial poisons
 - insecticides
 - herbicides
 - disinfectants such as triclosan
 - chemotherapy:
 - competitive inhibitors (Sulfonamides)
 - competition at the active site (treatment of methanol intoxication)
 - irreversible inhibition (nonsteroidal anti-inflammatory drugs –aspirin)
 - Anti-cancer drugs (Methotrexate – analogue of folic acid)

Principle serum enzymes used in clinical diagnosis

Serum Enzyme	Major Diagnostic Use
Aminotransferases	Myocardial infarction Viral hepatitis
Aspartate aminotransferase	
Alanine aminotransferase	
Amylase	Acute pancreatitis
Ceruloplasmin	Hepatolepticular degeneration
Creatine kinase	Muscle disorders and myocardial infarction
γ-Glutamyl transpeptidase	Various liver diseases
Lactate dehydrogenase (isozymes)	Myocardial infarction
Lipase	Acute pancreatitis
Phosphatase, acid	Metastatic carcinoma of the prostate
Phosphatase, alkaline (isozymes)	Various bone disorders, obstructive liver diseases

Introduction to nucleic acids

Role of nucleotides:

- the energy currency in metabolic transactions (ATP, GTP),
- the essential chemical participants in signals transduction paths of hormones and other extracellular stimuli in the of cells
- the structural components of enzyme cofactors and metabolic intermediates
- the molecular repositories of genetic information.

Gene is a segment of a DNA molecule that contains the information required for the synthesis of a functional biological product, whether protein or RNA.


Ribosomal RNAs (rRNAs) are components of ribosomes

Messenger RNAs (mRNAs) are intermediaries, carrying genetic information from one or a few genes to a ribosome



Transfer RNAs (tRNAs) are adapter molecules that faithfully translate the information in mRNA into a specific sequence of amino acids.

Structure of nucleotides

phosphoric ester bond



(a) nitrogenous (nitrogen-containing) base




(b) Pyrimidine Purine


nucleoside

Base	Nucleoside	Nucleotide	Nucleic acid
Purines			
Adenine	Adenosine Deoxyadenosine	Adenylate Deoxyadenylate	RNA DNA
Guanine	Guanosine Deoxyguanosine	Guanylate Deoxyguanylate	RNA DNA
Pyrimidines			
Cytosine	Cytidine Deoxycytidine	Cytidylate Deoxycytidylate	RNA DNA
Thymine	Thymidine or deoxythymidine	Thymidylate or deoxythymidylate	DNA
Uracil	Uridine	Uridylate	RNA

Major purine and pyrimidine bases of nucleic acids




Adenine




Guanine


Purines



Cytosine




Thymine
(DNA)



Uracil
(RNA)

Pyrimidines


Deoxyribonucleotides and ribonucleotides of nucleic acids



Nucleotide: Deoxyadenylate
(deoxyadenosine
5'-monophosphate)

Symbols: A, dA, dAMP


Nucleoside: Deoxyadenosine



Nucleotide: Deoxyguanylate
(deoxyguanosine
5'-monophosphate)

Symbols: G, dG, dGMP


Nucleoside: Deoxyguanosine



Nucleotide: Deoxythymidylate
(deoxythymidine
5'-monophosphate)

Symbols: T, dT, dTMP

Nucleoside: Deoxythymidine




Nucleotide: Deoxycytidylate
(deoxycytidine
5'-monophosphate)

Symbols: C, dC, dCMP

Nucleoside: Deoxycytidine


(a) Deoxyribonucleotides



Nucleotide: Adenylate (adenosine
5'-monophosphate)

Symbols: A, AMP


Nucleoside: Adenosine



Nucleotide: Guanylate (guanosine
5'-monophosphate)

Symbols: G, GMP


Nucleoside: Guanosine



Nucleotide: Uridylate (uridine
5'-monophosphate)

Symbols: U, UMP

Nucleoside: Uridine



Nucleotide: Cytidylate (cytidine
5'-monophosphate)

Symbols: C, CMP


Nucleoside: Cytidine

(b) Ribonucleotides


The nucleotide units of DNA (a) are usually symbolized as A, G, T, and C, sometimes as dA, dG, dT, and dC; those of RNA (b) as A, G, U, and C..

In their free form the deoxyribonucleotides are commonly abbreviated dAMP, dGMP, dTMP, and dCMP; the ribonucleotides, AMP, GMP, UMP, and CMP.


Some minor purine and pyrimidine bases




5-Methylcytidine




N⁶-Methyladenosine




(a) N²-Methylguanosine




5-Hydroxymethylcytidine




Inosine



Pseudouridine




(b) 7-Methylguanosine




4-Thiouridine


Some adenosine monophosphates




Adenosine 5'-monophosphate



Adenosine 2'-monophosphate



Adenosine 3'-monophosphate



Adenosine 2',3'-cyclic monophosphate



3,5-cyclic monophosphate (cAMP)
guanosine 3,5-cyclic monophosphate (cGMP)

Phosphodiester linkages in the covalent backbone of DNA and RNA

oligonucleotide


The backbone of alternating pentose and phosphate groups in both types of nucleic acid is highly polar. The 5 end of the macromolecule lacks a nucleotide at the 5 position, and the 3 end lacks a nucleotide at the 3 position.

5-phosphate group of one nucleotide unit is joined to the 3-hydroxyl group of the next nucleotide, creating a phosphodiester linkage



Some simpler representations of this pentadeoxyribonucleotide are:
pA-C-G-T-AOH, pApCpGpTpA, and pACGTA

Hydrogen-bonding patterns in the base pairs defined by Watson and Crick



- Stacking interactions:**
- *hydrophobic*
 - *van der Waals*
 - *dipole-dipole interactions*

Hydrogen bonds involving the amino and carbonyl groups



James D. Watson



Francis Crick,