

International higher school of medicine

Topic : pancreatic cancer

Done by : shantanu

Shreyash

shrinit

- Pancreatic cancer occurs when changes (mutations) in the pancreas cells lead them to multiply out of control. A mass of tissue can result. Sometimes, this mass is benign (not cancerous). In pancreatic cancer, however, the mass is malignant (cancerous).

- There are two types of tumors that grow in the pancreas: exocrine or neuroendocrine tumors. About 93% of all pancreatic tumors are exocrine tumors, and the most common kind of pancreatic cancer is called [adenocarcinoma](#). Pancreatic adenocarcinoma is what people usually mean when they say they have pancreatic cancer. The most common type begins in the ducts of the pancreas and is called ductal adenocarcinoma.
- The rest of the pancreatic tumors — about 7% of the total — are neuroendocrine tumors (NETs), also called pancreatic NETs (PNETs), an islet cell tumor or islet cell carcinoma. Some NETs produce excessive hormones. They may be called names based on the type of hormone the cell makes — for instance, insulinoma would be a tumor in a cell that makes insulin.

Incidence (2014 Data)

46,420 (2.8%)

Deaths

39,590 (6.8%)

The 5 year survival is only 6.7%

Life time risk of developing this cancer is 1.5%



2014 Statistics

Male

Female

New Cases 23,530 (2.7%) 22,890 (2.8%)

Deaths 20,170 (6.5%) 19,420 (7%)


Risk factors for pancreatic cancer include the following

- Smoking
- Obesity
- Personal history of diabetes or chronic pancreatitis
- Family history of pancreatic cancer or pancreatitis
- Certain hereditary conditions


Symptoms

- Most patients present with pain (in the back) weight loss or jaundice
- Tumors in the head of the pancreas are more likely to have jaundice,
- Those that arose in the body or tail, more likely pain and weight loss.


Bile: yellowish fluid produced in the liver that aids in digestion of fat in the small intestine




Bile: passes through the common bile duct through the head of the pancreas on it's way to the duodenum




Bile duct carries the bilirubin through the head of the pancreas on it's way to the duodenum




Tumors in the head of the pancreas are more likely to have jaundice




Tumors in the head of the pancreas are more likely to have jaundice





Tumors in the body or tail are more likely to present with pain or weight loss



Symptoms of Pancreas Cancer

- Asthenia (weakness) 86 percent
- Weight loss 85 percent
- Anorexia (no appetite)-83 percent
- Abdominal pain - 79 percent Epigastric pain (stomach)-71 percent • Dark urine-59 percent
- Jaundice 56 percent
- Nausea 51 percent Back pain - 49 percent
- Diarrhea- 44 percent
- Vomiting-33 percent
- Steatorrhea (fatty stools)- 25 percent
- Thrombophlebitis - 3 percent


Signs of Pancreas Cancer

- Jaundice (yellow) - 55 percent
- Hepatomegaly (large liver) - 39 percent
- Right upper quadrant mass - 15 percent
- Cachexia (wasting) 13 percent
- Courvoisier's sign (nontender but palpable distended gallbladder at the right costal margin) 13 percent
- Epigastric mass (felt lump in stomach) - 9 percent
- Ascites (abdominal fluid) - 5 percent


Pathology

- Ductal adenocarcinoma accounts for about 85% of all neoplasms. And more than 95% of all pancreatic cancers arise from the exocrine (digestive enzymes) elements. Cancers that arise from the endocrine cells (neuroendocrine, islet cells) account for 5% or less

Part of pancreas



Cancer in the Head of the Pancreas



Location of Pancreas Cancer

- 60 to 70 percent of exocrine pancreatic cancers are localized to the head
- 20 to 25 percent are in the body/tail and the remainder involve the whole organ
- H = Head
- N = Neck
- B = Body
- T = TailUn Uncinate

Tests used to evaluate and stage pancreas cancer

- Routine blood tests e.g. liver products like bilirubin
- Elevated tumor markers (CA 19-9 or CEA)
- MRI, CT scans, Ultrasound
- Endoscopy including endoscopic ultrasound or ERCP
- Laparoscopy Biopsy

CA 19-9

- The reported sensitivity and specificity rates of CA 19-9 for pancreatic cancer range from 70 to 92, and 68 to 92 percent, respectively. The rates of unresectable disease among all patients with a CA 19-9 level ≥ 130 units/mL versus < 130 units/mL were 26 and 11 percent, respectively. Among patients with tumors in the body/tail of the pancreas, more than one-third of those who had a CA 19-9 level ≥ 130 units/mL had unresectable disease.

Elevated CA 19-9

- Cancer


1. Pancreas
2. Biliary Cancer (gallbladder, cholangiocarcinoma, ampullary)-
Hepatocellular
3. Gastric, ovarian, colorectal (less often)
4. Lung, breast, uterus (rare)

- Benign

- Acute cholangitis
- Cirrhosis and other cholestatic diseases (gall stones)

Ultrasound

- study of 900 patients who underwent ultrasound to work up painless jaundice, anorexia, or unexplained weight loss. The sensitivity for detection of all tumors in the pancreas was 89 percent. Among the 779 patients who were followed over time and established not to have developed a pancreatic tumor, nine had false-positive US findings (specificity 99 percent).




Mass in the pancreatic head and dilated common bile duct and pancreatic duct


CT

- Sensitivity of CT for pancreatic cancer depends on technique and is highest (89 to 97 percent) with triple-phase, helical multidetector row CTAs expected, sensitivity is higher for larger tumors; in one study, the sensitivity was 100 percent for tumors >2 cm, but only 77 percent for tumors ≤ 2 cm in size

Endoscopy or ERCP or EUS




ERCP or Endoscopic retrograde cholangiopancreatography



ERCP

Try to
get into
the duct
system





ERCP
Inject dye into
the duct system
and look for
compression





ERCP Inject dye into the duct system and look for compression


ERCP

- Sensitivity of 92 percent and Specificity of 96 percent for diagnosing cancer of the pancreas by ERCP. ERCP provides an opportunity to collect tissue samples (forceps biopsy, brush cytology) for histologic diagnosis. However, the sensitivity for detection of malignancy (approximately 50 to 60 percent) is lower than that of endoscopic ultrasound (EUS)-guided FNA (sensitivity 92 percent)


Endoscopic Placement of a Stent




CT Scan - Pancreas Cancer




CT Scans




CT or PET Scan







PET Scans




Stage IA (T1aN0M0)




Stage IB (T2N0M0) over 2cm, limited to pancreas




Stage IIA (T3N0) beyond the pancreas



Stage IIB (T1-3N1M0)



- **Stage III (T4) Unresectable Cancer** has spread to the major blood vessels near the pancreas. These include the superior mesenteric artery, celiac axis, common hepatic artery, and portal vein.



Thank you