



OPHTHALMOLOGY: OUR RESOURCES






UP-TO-DATE DIAGNOSTIC METHODS

OPTICAL COHERENCE TOMOGRAPHY



Stratus OCT



Non-invasive optical diagnostic techniques that would provide images of high resolution for the measurement and diagnosis of ocular tissues, it useful for management and follow-up of eye pathology



OPTICAL COHERENCE TOMOGRAPHY OF ANTERIOR EYE SEGMENT


○ *Visante OCT*





OCT OF ANTERIOR EYE SEGMENT

- provides high-resolution anterior chamber images;
- measurement of anterior segment ocular structures, including anterior chamber depth (ACD), anterior chamber angles, corneal thickness
- allows to evaluate the potential risk of glaucoma, complications for cataract surgery



THE NON-MYDRIATIC FUNDUS CAMERA SYSTEM FROM CARL ZEISS



TF



- Brilliant true color retinal images
- Automontage for panoramic retinal overviews
- Documentation of the results and possibility of subsequent management of the disease

**The Non-Mydriatic
Fundus Camera**




Ultrasound eye examination





Fluorescein angiography





OPHTHALMOSURGERY



OPERATING- ROOM







HIGHLY SKILLED OPHTHALMOSURGEONS





PHACOEMULSIFICATION CATARACT SURGERY

- Microincision (less 2,4 mm) – without sutures;
- Decrease of post-operative astigmatism
- Decrease of possible intra- and post-operative complications





Premium-class phacomashines







Infiniti





Accurus

PHACOEMULSIFICATION OF DIFFERENT TYPES OF CATARACT





PHACOEMULSIFICATION CATARACT «WITH WATER»





IOL-MASTER FOR IOL POWER CALCULATION




Quickly and accurately!



EXCELLENT VISION AT MULTIPLE DISTANCES

Acrysof IQ ReSTOR +3,0D:

- vision at near, intermediate and distance
- **not needing glasses**




**AcrySof® ReSTOR®
ASPHERIC +3**




TORIC IOL FOR PRECISE ASTIGMATISM CORRECTION



mark the cornea prior to
implantation







Toric IOL

◎ New quality of vision












MARK THE CORNEA PRIOR TO IMPLANTATION



With the patient sitting upright and looking at a distance target, the cornea is marked at the 3:00 and 9:00 positions

IN OPERATING-ROOM










IMPLANTATION OF IOL WITH SULCUS FIXATION


In cases:

- after eye trauma
- in post-operative aphakia
- in zonulla thickness



VITREO-RETINAL SURGERY









Accurus




BIOM



VITRECTOMY WITH ENDOLASERCOAGULATION



До операции



После операции

Before surgery



After Surgery








CORNEAL TRANSPLANTATION




- Keratoconus
- Endothelial-epithelial corneal dystrophy
 - scarring due to keratitis



Penetrating keratoplasty



Before surgery



After surgery