

Research in Medicine:

Basic Types and Methods



Medical Research and Study Design

- The scientific value and informativeness of a medical study are determined to a major extent by the study design.
- Errors in study design cannot be corrected afterwards.

5 phases of medical research

- Planning
- Performance
- Documentation
- Analysis
- Publication



The importance of study design

Planning  study design:

a suitable type of study

an overall plan for all procedures

a bias to be minimized

6 aspects of study design

1. the question to be answered,
2. the study population,
3. the type of study,
4. the unit of analysis,
5. the measuring technique
6. the calculation of sample size

The question to be answered

- research objectives = the key research question must be
- operationalized → converted into a measurable and evaluable form.
- main questions vs secondary questions.
- as a result open questions are answered and new hypotheses are possibly generated.

Descriptive analysis

the units of analysis are to be described by the recorded variables (e.g. blood parameters or diagnosis)




Exploratory analysis

- to recognize connections between variables,
- to evaluate connections,
- to formulate new hypotheses.



Confirmatory analysis

to provide statistical proofs by testing specified study hypotheses



Unit of analysis

clinical study

- patient
- hereditary information,
- a cell,
- a cellular structure,
- an organ, an organ system,
- a single test individual (animal or man),
- a specified subgroup or
- the population of a region or of a country.


systematic reviews

- a single study
- The interesting information or data (observations, variables, characteristics) are recorded for the statistical units.

Measuring technique

measuring *instruments*:


- to record measuring data (such as blood pressure or laboratory parameters)
- to collect data with standardized or self-designed questionnaires (for example, quality of life, depression, or satisfaction).



Classification of study types


Primary Research

actual studies



Secondary Research

summarizes available studies in the form of reviews and meta-analyses



Primary Research: Main Categories

- basic medical research (or experimental research)
- clinical research
- epidemiological research

BASIC MEDICAL RESEARCH



also known as **experimental** research

Aim: to **acquire new knowledge** without looking for long-term benefits other than the advancement of knowledge

includes: animal experiments, cell studies, biochemical, genetic and physiological investigations, and studies on the properties of drugs and materials



CLINICAL RESEARCH

- **interventional (or experimental)**
studies on drugs, medical devices and
studies in which surgical, physical or
psychotherapeutic
procedures are examined.

- **noninterventional (or observational)** -
a study in which knowledge from
the treatment of persons is
analyzed using epidemiological
methods. The diagnosis,
treatment and monitoring are
performed exclusively
according to medical practice.

Clinical Studies (Clinical Trials)

Aim: to determine better ways to prevent, screen for, diagnose or treat diseases.

Controlled trial (a study group and a control group)

Randomization (a randomized controlled trial)

Blinding (single and double blind trial)

EPIDEMIOLOGICAL RESEARCH



the description of *health and welfare* in populations through the collection of data related to health and the frequency, distribution and determinants of disease in populations, with the aim of improving health.

Epidemiological: interventional or experimental

field studies

- sample from an area, such as a large region or a country

e.g. investigation of the iodine supplementation of cooking salt to prevent cretinism in a region with iodine deficiency

group studies

- sample from a specific group, such as a specific social or ethnic group.

Epidemiological: observational

- Case control studies


Cases are compared with controls. Cases are persons who fall ill from the disease in question. Controls are persons who are not ill, but are otherwise comparable to the cases. A retrospective analysis is performed to establish to what extent persons in the case and control groups were exposed. Possible exposure factors include smoking, nutrition and pollutant load.

Epidemiological: observational


- Cohort studies (prospective and retrospective)

cohort studies involve the observation of **two healthy groups of subjects** over time. One group is exposed to a specific substance (for example, workers in a chemical factory) and the other is not exposed. It is recorded prospectively (into the future) how often a specific disease (such as lung cancer) occurs in the two groups.

a



b



Common study types for epidemiological investigations

Study objective

Study of severe diseases
such as cancers

Study of rare exposure,
such as exposure to
industrial chemicals

Study of multiple
exposures, such as the
combined effect of oral
contraceptives and
smoking on myocardial
infarction

Study type

Case control studies

Cohort studies in a
population group (e.g.
industrial workers

Case control studies

(Dtsch Arztebl Int. 2009 April;
106(15): 262–268.)

References

- [http:// www.experiment-resources](http://www.experiment-resources)
- <http://www.nhs.uk/Conditions/Clinical-trials>
- Glossary of Clinical Trials Terms
- Altman DG. Randomisation: potential for reducing bias. BMJ. 1991;302:1481–1482.
[PMC free article] [PubMed]
- Study Design in Medical Research // Dtsch Arztebl Int. Mar 2009; 106(11): 184–189.
Published online Mar 13, 2009. doi: [10.3238/arztebl.2009.0184](https://doi.org/10.3238/arztebl.2009.0184)
- Types of Study in Medical Research // Dtsch Arztebl Int. Apr 2009; 106(15):
262–268. Published online Apr 10, 2009. doi: [10.3238/arztebl.2009.0262](https://doi.org/10.3238/arztebl.2009.0262)
- Study designs in medical research <http://galton.uchicago.edu/>

Thank you for attention

