




**Молекулярная генетика развития**  
**Вельков В В 2013**  
**Gene Regulation during Development**

# Morphogenesis - How do you get from a spherical egg to say a frog?



# Dolly and Bonnie




The nucleus from an differentiated adult cell (from the mammary gland) was able to replace the nucleus of a fertilized egg and produce an adult complete with functional gametes: Dolly had a daughter – Bonnie. SO: during development genes are just turned on and off in a carefully orchestrated fashion. SO, THE DOGMA OF DEV. BIOL. Is differential gene expression controls development



The Spemann Experiment  
Spemann & Mangold, 1923

# Spemann Experiment

- Outline: Graft a tissue from one embryo into another embryo, and see what happens – **Cut and Paste**




- **Main observation: A graft of a specific tissue (the organizer) to a specific location can induce Siamese twins connected at the belly.**
- **Conclusion: Hilde Mangold and Hans Spemann found that a piece of Prospective dorsal tissue can organize formation of an embryo**



“A piece taken from the upper blastopore lip of a gastrulating amphibian embryo **exerts an organizing effect on its environment** ...  
Such a piece can therefore be designated as an **organizer**”

Original blastopore lip

Graft



Vegetal view






- Spemann won the Nobel prize in 1935
- Hilde Mangold died in 1926...
- “Spemann Organizer” was found in all vertebrates, including human

The organizer secretes both **morphogens** and their inhibitors which diffuse throughout the embryo

# Universal mechanism of animal development




Short- long-range diffusible signaling molecule


**Gene expression controls 4 essential process**

# Development is progressive

- **Specification of cell fate: determination**
  - All cells still ‘look the same’
  - Can be tested by transplantation experiments
- **Interactions can make cells different from each other: induction**



- In the process called induction
  - Signal molecules from embryonic cells cause transcriptional changes in nearby target cells




**(b) Induction by nearby cells.** The cells at the bottom of the early embryo depicted here are releasing chemicals that signal nearby cells to change their gene expression.

# From single cell to organism – a life cycle

- Fertilisation followed by cell division
- Pattern formation – instructions for
  - Body plan (Axes: A-P, D-V)
  - Germ layers (ecto-, meso-, endoderm)
- Cell movement - form – gastrulation
- Cell differentiation
- Cell growth, cell death (apoptosis)


The use of a model organism





# Three Strategies:

- mRNA localization
- Cell-to-cell contact
- Signaling through the diffusion of secreted signaling molecules



A movie poster for the film 'The Meeting Place Cannot Be Changed'. The image shows two men in a war-torn city. The man on the left has a mustache and is wearing a dark uniform. The man on the right is wearing a dark jacket with a white star on the lapel. The background shows damaged buildings and a red, hazy sky. The title is written in large, bold, red Cyrillic letters at the bottom.

**МЕСТО  
ВСТРЕЧИ  
ИЗМЕНИТЬ  
НЕЛЬЗЯ**



Морфогены  
и рецепторы  
морфогенов



**МЕСТО  
ВСТРЕЧИ  
ИЗМЕНИТЬ  
НЕЛЬЗЯ**


- **Morphogen – substances that define different cell fates in a concentration-dependent manner**
- **Interaction of two signaling centers located in the anterior and posterior poles of the egg pattern insect body axis**

- **Morphogen** = Soluble molecule that causes cellular commitment but is secreted some distance from the target cells.
- **Morphogen Gradient** = concentration


# Градиенты морфогенов

- Белки, кодируемые генами, функционирующими в ходе созревания яйца, и транспортируемые туда из питающих клеток,
- распределяются по оси яйца,
- образуя градиенты, специфически характерные для продуктов каждого гена


# Morphogen Threshold Concentrations



# Morphogen Threshold Concentrations



# Morphogen Threshold Concentrations



# Morphogen Receptor

Some mRNAs Become Localized within Eggs and Embryos due to an Intrinsic Polarity in the Cytoskeleton

- The asymmetrically distributed mRNA is transported along element of the cytoskeleton
- from – to the growing + end.



## Adaptor protein

- The **morphogene receptor** binds to 3'untranslated trailer (3'UTR) region

**of morphogene mRNA,**

A morphogene receptor


has two domains:

one recognizes the 3' UTR,  
the other

associates with myosin.

Cytoskeletal elements –

microtubules are polarized within the oocyte and can be used to allow the localization of mRNA molecules to specific parts of the cell.



# Influence of Other Cells

- **Morphogen Receptor Gradient** = frequency gradient of the receptors for a morphogen in target cell cell membranes




Morphogen gradient

The diagram consists of a vertical rectangular box with a teal border. The box is divided into two horizontal sections. The top section is shaded in a light blue gradient, and the bottom section is shaded in a darker blue gradient. The text 'Morphogen gradient' is centered in the top section, and 'Morphogen receptor gradient' is centered in the bottom section.

Morphogen receptor gradient


# Morphogens

- Coordinate the cell growth and differentiation.
- Formation of a **long-range concentration profile**.
- Cellular response is concentration dependent.






# Life cycle




# Many Adult Structures Develop from Imaginal Discs in Larvae and Pupae




# Position or Spatial Organization is *Everything*

- 2 main mechanisms used to communicate positional information
  1. Morphogens
  2. Cell adhesion




Fate Map: all **different regions** of an embryo develop is described

Developmental program — **process** for cell achieving its fate







**(a) The first three hours after fertilization**




**(b) Early embryonic stages in cross section**




(a) Cellular blastoderm




(b) Gastrulation



(c) Segmentation




(d) Segment identity is preserved throughout development.



- **Морфогены активируют**
- **Гены сегментации активируют**
- **Гомеозисные гены активируют...**

# ***Genetic Analysis of Early Development: Scientific Inquiry***

- **The study of developmental mutants**
  - **Laid the groundwork for understanding the mechanisms of development**



Wild type

Mutant

# *Drosophila* Embryogenesis

First phase is establishment of body axes

**Before** fertilization, specialized **nurse cells**


move maternal mRNAs (morphogenes) into maturing oocyte

Then morphogens are distributed in the oocyte prior to fertilization






- **The Morphogens organize the structure of the egg**

These positioned mRNA will initiate a cascade of gene activations following fertilization

- Яйцо созревает в особой камере — фолликуле.
- В фолликулах - гены с **материнским эффектом**, которые функционируют еще до оплодотворения яйца сперматозоидом
- В фолликуле ооцит и 15 огромных питающих клеток,
- которые синтезируют «РНК морфогены» и перекачивают их в ооцит.



At the start of development, **gradients** are established in the egg along two axes, **anterior-posterior** and **dorsal-ventral**.



**b.**

**e**

## Anterior determination

Egg

*bicoid* RNA



- 1 *bicoid* RNA is localized in the anterior region of the egg.

*bicoid* protein



- 2 *bicoid* protein forms a gradient by diffusing toward the posterior of the embryo.

*hunchback* protein



- 3 *hunchback* is transcribed and translated in the anterior of the embryo.

1. Maternal gene products, called **morphogens**, establish gradients in early embryogenesis.
2. Anterior-posterior development uses localized gene regulators.
3. Dorsal-ventral development uses localized receptor-ligand interactions.


Syncytial blastoderm

Cellular blastoderm



# Градиент морфогенов активирует зиготные гены

- После того как градиенты в яйце созданы, происходит оплодотворение и начинается дробление зародыша
- образуется однослойная бластодерма. Каждая клетка в ней занимает определенное положение по отношению к сформировавшимся градиентам,
- - обладает определенной позиционной информацией.
- Морфогены взаимодействуют с регуляторными участками генов, активирующихся у зигот (зиготических генов).



**Expressed during oogenesis by the mother.** Act upon or within the maturing oocyte.

**Expressed after fertilization.** Mutations in these genes alter the number or polarity of segments. Three groups of segmentation genes act sequentially to define increasingly smaller regions of the embryo.

**Control the identity of a segment,** but do not affect the number, polarity or size of segments. Mutations in these genes cause one body part to develop the phenotype of another part.

# Establishment of the A/P axis

Nurse cells secrete maternally produced *bicoid* and *nanos* mRNAs into the oocyte



- Differentially transported by microtubules to opposite poles of the oocyte

  - bicoid* mRNA to the future anterior pole

  - nanos* mRNA to the future posterior pole


- After fertilization, translation will create opposing gradients

of **Bicoid** and **Nanos** proteins



# *bicoid* (*bcd*) Encodes the Anterior Morphogen

(a) Localization of *bicoid* mRNA



Anterior

Posterior

(c) Bicoid protein is a morphogen.

Mother

*bicoid*<sup>+</sup>/*bicoid*<sup>-</sup>

1 dose

*bicoid*<sup>+</sup>/*bicoid*<sup>+</sup>


2 doses

*bicoid*<sup>+</sup>


4 doses

*bicoid*<sup>+</sup>

6 doses




# Bicoid gradient



- **Bicoid acts as a transcriptional activator of *hunchback* gene transcription**

# Nanos gradient



# Establishment of the A/P axis


Bicoid and Nanos control translation of two other maternal mRNAs, *hunchback* and *caudal*, that encode transcription factors

- Hunchback** activates anterior structures
- Caudal** activates posterior structures


The two mRNAs are not evenly distributed

- Bicoid protein inhibits *caudal* mRNA translation
- Nanos protein inhibits *hunchback* mRNA translation






**a. Oocyte mRNAs**




**b. After fertilization**




**c. Early cleavage embryo proteins**

# How Bcd Protein Works



## Proteins in early cleavage embryos



# Establishment of the D/V axis

Maternally produced **dorsal mRNA** is placed into the oocyte

- Not asymmetrically localized


Oocyte nucleus synthesizes *gurken* mRNA

- Accumulates in a crescent on the future dorsal side of embryo

After fertilization, a series of steps results in selected transport of **Dorsal** into ventral nuclei, thus forming a D/V gradient


# Segmentation genes act sequentially to divide the embryo into segments

- Normal *Drosophila* embryo divided into 15 segments
  - 3 head, 3 thoracic and 9 abdominal
  - Each will give rise to unique morphological features in adult



Парасегменты (compartments) соответствуют границам экспрессии различных генов сегментации и смещены влево от морфологически выраженных сегментов (на пол сегмента)

## Segmentation in Drosophila



(Klug & Cummings 1997)

# Production of Body Plan

The body plan is produced by sequential activation of three classes of **segmentation genes**

Segmentation Is Initiated by Localized RNAs at the Anterior and Posterior Poles of the Unfertilized Egg

## 1. Gap genes

- Map out the coarsest subdivision along the A/P axis
- All 9 genes encode transcription factors that activate the next gene class


## 2. Pair-rule genes

- Divide the embryo into seven zones
- The 8 or more genes encode transcription factors that regulate each other, and activate the next gene class

## 3. Segment polarity genes

- Finish defining the embryonic segments

# 3 группы генов сегментации




- *Gap-гены,*
- *гены pair-rule*
- *гены сегментной полярности.*



- **Gap genes**
  - **Expressed first**
  - **Gap mutants show a gap in segmentation pattern at positions where particular gene is absent**
  - **Binding sites in promoter have different affinities for maternal transcription factors**
  - **Gap genes encode transcription factors that influence expression of other gap genes**


# Gap genes



- *Hunchback*, *krüppel*, *giant*, *tailless* and *knirps*.
- Their expression patterns in the early embryo are determined by the maternal effect gene products
- **These genes establish the segmented body plan of the embryo along the anterior-posterior axis.**



# The A-P axis is divided into broad regions by gap gene expression

- The first *zygotic* genes
- Respond to maternally-derived instructions
- Short-lived proteins,
- gives bell-shaped distribution from source




# Zones of Expression of Four Gap Genes: *Hunchback*, *Krüppel*, *Knirps*, and *Giant* in Late Syncytial Blastoderm Embryos


(a) Zones of gap gene expression



# Defects in Segmentation from Mutations in Gap Genes





# Mutations in Gap Gene Result in Loss of Segments Corresponding to Zone of Expression



## Transcription factors in cascade

# GAP genes



- ***Hunchback (hb)***, a gap gene, responds to the dose of bicoid protein
- A concentration above threshold of bicoid activates the expression of *hb*
- The more *bicoid* transcripts, the further back *hb* expression goes



## *Krüppel* reads two values

### GAP genes

- *Krüppel* (*Kr*), a gap gene, responds to the dose of hb protein
- A concentration **above minimum threshold** of hb activates the expression of *Kr*
- A concentration **above maximum threshold** of hb inactivates the expression of *Kr*







## Establishing the Polarity of the Embryo

Fertilization of the egg triggers the production of bicoid protein from maternal RNA in the egg. The bicoid protein diffuses through the egg, forming a gradient. This gradient determines the polarity of the embryo, with the head and thorax developing in the zone of high concentration (green fluorescent dye in antibodies that bind bicoid protein allows visualization of the gradient).



## Setting the Stage for Segmentation

About 2 1/2 hours after fertilization, bicoid protein turns on a series of brief signals from so-called gap genes. The gap proteins act to divide the embryo into large blocks. In this photo, fluorescent dyes in antibodies that bind to the gap proteins Krüppel (orange) and Hunchback (green) make the blocks visible; the region of overlap is yellow.





## Laying Down the Fundamental Regions

About 9.5 hr later, the gap genes switch on the "pair-rule" genes, which are each expressed in seven stripes. This is shown for the pair-rule gene *hairy*. Some pair-rule genes are only required for even-numbered segments while others are only required for odd numbered segments.



## Forming the Segments

The final stage of segmentation occurs when a "segment-polarity" gene called engrailed divides each of the seven regions into halves, producing 14 narrow compartments. Each compartment corresponds to one segment of the future body. There are three head segments (H, bottom right), three thoracic segments (T, upper right), and eight abdominal segments (A, from top right to bottom left).



We know some mysterious mutations,  
which generate horroristic monsters



**Homeotic  
mutations**

# Homeotic-Selector/ HOX Genes

# Production of Body Plan

Segment identity arises from the action of **homeotic genes**

-Mutations in them lead to the appearance of normal body parts in unusual places


-*Ultrabithorax* mutants produce an extra pair of wings



# Production of Body Plan


## Homeotic gene complexes

- The **HOM complex** genes of *Drosophila* are grouped into two clusters
  - Antennapedia complex**, which governs the anterior end of the fly
  - Bithorax complex**, which governs the posterior end of the fly
- Interestingly, the order of genes mirrors the order of the body parts they control***



**Homeotic genes specify body segment identity in *Drosophila*.**

# Homeotic Genes Determine Specialization of Segments



# Гомеозисные гены

- Большой класс генов
- контролируют развитие части тела
- из определенного сегмента.
  
- В результате гомеозисной мутации
- из данного сегмента развивается
- другая часть тела


Наиболее известны

- *Antennapedia-complex (Ant-C).*
- *Bithorax-complex (BX-C)* и




- *Antennapedia complex*
- 5 genes that affect the anterior part of the fly
- When mutated, legs grow instead of antennae

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




(a) Normal fly




(b) Mutant fly

a: © Juergen Berger/Photo Researchers, Inc.; b: F. R. Turner, Indiana University


# Homeotic Genes: Master Regulatory Genes




# Homeotic mutation: Master Regulatory Genes



Normal fly



*Antennapedia* mutant fly





# Halteres into wings



One structure is placed in the position where normally another structure would be (e.g. a pair of wings instead of halteres)

Gilbert, SF (2003) Developmental Biology, 7<sup>th</sup> ed.

Гомеозисная мутация у *Drosophila* (подавление активности гена *ultrabithorax*) приводит к образованию второй пары крыльев вместо жужжалец



wildtype





Ultrabithorax

Гомеозисные гены относятся к селекторным генам. Через свои белковые продукты -транскрипционные факторы - они избирательно воздействуют на гены начальных звеньев генетических формообразовательных программ, определяя выбор дифференцировки целого участка тела организма. Кодированные ими транскрипционные факторы для избирательности воздействия могут использовать кофакторы. У *Drosophila* - это белковые продукты генов *Extradenticle*, *tearshirt* и др.

## Гомеозисные (гомеотические) гены

Активности гомеозисных генов иницируются продуктами экспрессии генов сегментации, а затем поддерживаются самостоятельно



**Figure 9.8**

Generalized model of *Drosophila* anterior-posterior pattern formation. (A) The pattern is established by maternal effect genes that form gradients and regions of morphogenetic proteins. These morphogenetic determinants create a gradient of Bicoid protein that differentially activates the gap genes, which define broad territories of the embryo. The gap genes enable the expression of the pair-rule genes, each of which divides the embryo into regions about two segments wide. The segment polarity genes then divide the embryo into approximately 100 units along the anterior-posterior axis. Together the actions of these genes define the spatial domains of the homeotic genes that define the identities of each of the segments. In this way, periodicity is generated from nonperiodicity, and each segment is given a unique identity. (B) Maternal effect genes. The anterior axis is specified by the gradient of Bicoid protein (yellow through red). (C) Gap gene protein expression and overlap. The domains of Bicoid protein (orange) and the domains of Krüppel protein (green) overlap to form a region containing both transcription factors (yellow). (D) Products of the pair-rule genes form seven bands across the embryo. (E) Products of the segment polarity genes expressed, seen here at the extended germ band stage. (B courtesy of C. Nüsslein-Volhard; C courtesy of C. Tsao and M. Levine; D courtesy of T. Korn; E courtesy of S. Carroll and S. Paddock.)



# Production of Body Plan

## Homeotic gene complexes


- All of these genes contain a conserved 180-base sequence, the **homeobox**
- Encodes a 60-amino acid DNA-binding domain, the **homeodomain**
- Homeobox-containing genes are termed ***Hox* genes**
- Vertebrates have 4 *Hox* gene clusters

# Production of Body Plan

## *Drosophila* HOM

### Chromosomes

#### *Drosophila* HOM genes





Fruit fly embryo



Fruit fly

## Mouse Hox

### Chromosomes



Mouse embryo




Mouse

a

b.

# Как произошли мухи?




Мухи эволюционировали из насекомых, имевших 4 крыла, Насекомые произошли из членистоногих, имевших множество ног. В ходе эволюции у мух сформировались группы генов: а) подавляющие развитие ног на брюшных сегментах многоножкоподобных предков, б) подавляющих развитие второй пары крыльев, в) формирующие новые структуры: гальтеры и брюшные сегменты.

# К чему приводит делеция *VX-C*?

- Эмбрион развивается до определенной стадии и затем гибнет
- Эмбрион имел
  - проторакальный сегмент, а
  - все остальные — мезоторакальные
- Если бы этот организм выжил до взрослой мухи,
  - она бы имела 10 пар крыльев и 10 пар ног
- **Функция гена *VX-C***
  - - инактивация генов, формирующих ноги и крылья во всех последующих после второго торакального сегментах,
  - - развитию всех структур на брюшных сегментах

Предполагаемая макроэволюционная роль гомеозисных мутаций - вклад биологии развития в эволюционную теорию. Однократная мутация одного или группы *Ном-С* генов может привести к образованию (утрате) пары крыльев или ножек




**Figure 22.4**

Differences in larval and adult morphology due to Hox gene differences. (A, B) Larva and adult of *Drosophila*, a dipteran. An arrow points to one of the halteres of the adult. The larva lacks prolegs; its anterior end is at the left. (C, D) Larva and adult of *Danaus plexippus*, the monarch butterfly, a lepidopteran. The anterior of the caterpillar is to the left, and a proleg is indicated by an arrow. The adult has hindwings rather than halteres. The regulation of Hox genes determines the presence of prolegs, and the targets of Hox genes determine whether the third thoracic segment is to generate halteres or hindwings. (A courtesy of M. Tyler; B courtesy of E. B. Lewis; C courtesy of G. Savage; D by Bill Beatty/Visuals Unlimited.)

Кластер гомеозисных генов *Drosophila* - Hom-C. Активность генов в зародыше и во взрослой особи. Последовательное расположение генов в кластере (на 3-й хромосоме) определяет последовательность их экспрессии в парасегментах вдоль передне-задней оси

**Figure 9.27**

Homeotic gene expression in *Drosophila*. In the center are the genes of the Antennapedia and bithorax complexes and their functional domains. Below and above the gene map are the regions of homeotic gene expression (both mRNA and protein) in the blastoderm of the *Drosophila* embryo and the regions that form in the adult fly. Darker shaded areas are those segments or parasegments with the most product. (After Dessain et al. 1992 and Kaufman et al. 1990.)




**Vertebrates have four Hox complexes,  
with about 10 genes each.**

**They can be aligned in 13 groups of paralogues.**

**They display colinearity:**


- a) Temporal colinearity: genes on one end of the complex are expressed first, those on the other (posterior) end are turned on last.**
- b) Spatial colinearity: the more anteriorly expressed genes are in one end, the more posterior ones at the other end of the gene complex.**
- c) Anterior Hox genes are activated sequentially by retinoic acid.**

Районы экспрессии гомеотических генов в бластодерме зародыша *Drosophila* и несколькими часами позже в нервной системе (тёмные полосы - белок, светлые - иРНК)





# Temporal and spatial colinearity: order of Hox genes in DNA follows the antero-posterior body axis.




hindbrain and spinal cord


mesoderm

anterior


posterior



- **An identical or very similar nucleotide sequence**
  - **Has been discovered in the homeotic genes of both vertebrates and invertebrates**




**Кластеры гомеозисных генов высоко консервативны. Они сформированы в явном виде у Bilateria и присутствуют в организмах всех типов животных, включая человека.**






**Figure 3** Distribution of Hox genes in bilaterians. The tree on the left summarizes the 18S rDNA phylogeny of Aguinaldo *et al.*<sup>5</sup>. B, common bilaterian ancestor; D, common deuterostome ancestor; E, stem ecdysozoan; L, stem lophotrochozoan; P, common protostome ancestor. Horizontal black lines indicate mapping data (when available). Vertical white bars delineate orthologous genes or groups of


genes. Uncertain orthology relationships are indicated by question marks, dashed boxes indicate short PCR fragments, solid boxes indicate complete or almost complete homeoboxes, and striped colours indicate fast-evolving arthropod Hox sequences. The organisms studied in this paper are underlined.

У позвоночных выявляется не менее четырех кластеров паралогичных гомеозисных генов. Число генов в кластерах выше, чем у *Drosophila*, но не все паралогичные гены сохраняются в отдельных кластерах




**Figure 22.11**  
 Postulated ancestry of the homeotic genes from a hypothetical ancestor of both deuterostomes and protostomes. Amphioxus has only one cluster, similar to that of insects. Vertebrates have four clusters, none of which is complete. (After Holland and Garcia-Fernández 1996.)





# A summary of gene activity during *Drosophila* development





# Homeobox & Homeodomain

- ДНК связывающий модуль
  - факторов транскрипции
  - генов дифференцировки
- 
- 180 пн
  - 60 аминокислотных остатков




Гомеотические гены кодируют транскрипционные факторы -ТФ, содержащие ДНК- связывающий гомеодомен (60 а.о.), имеющий структуру - спираль-поворот-спираль- Helix-Turn-Helix (НТН). Участок ДНК, кодирующий гомеодомен - гомеобокс (180 н.п.). Не все гомеобокс-содержащие гены - гомеотические, а только входящие в состав кластера генов - Hom-C. Все гомеодомен-содержащие ТФ одинаково узнают последовательность ТААТ, но различно фланкирующие ТААТ нуклеотиды.









HOMEBOX PROTEIN VND / DNA COMPLEX (1NK2)




**HOMEODOMAIN PROTEINS** bind to DNA and regulate gene expression. They are composed of a variable region, which determines a protein's specific activity, a small connective hinge region and a homeodomain, a 60 amino acid sequence that is similar in all proteins of this type. This sequence is encoded by the homeobox regions of genes. The homeodomain consists of four alpha helices (1-4), one of which (*red*) recognizes and binds to a specific DNA sequence in the target genes.




**GRADIENTS** of homeodomain proteins can be seen in these stained limb buds. In the chick wing bud (*left*), the concentration of Hox 5.2 protein is highest in the cell nuclei near the posterior region (*right edge*). In the pectoral fin bud of a

zebrafish (*right*), the concentration of XlHbox 1 protein is highest in the anterior region (*left edge*). The fish pectoral fin is the evolutionary precursor of the tetrapod forelimb. Gradients are efficient mechanisms for conveying positional information.


# Hox Genes and EvoDevo




A= anterior  
 Group 3  
 C= Central  
 P=Posterior



**Mutation in HoxD13—synpolydactyly**  
**Extra digits & interphalangeal webbing (hetero)**  
**Similar but more severe & bony malformation**  
**of hands, wrists (Homo)**



**Loss of function-mild**  
**Gain of function (Polyalanine expansion**  
**in HoxD13) similar to multiple**  
**loss of function-HoxD11,12,13**



**Polyalanine expansion-**  
**Interfere all three proteins**  
**Semidominant effect**

## Effect of very high doses of retinoic acid on Hox gene expression in mouse embryos



**Normal mouse embryo**



**Retinoic acid:  
loss of many  
vertebrae**




**More retinoic acid:  
no posterior region  
formed**

**Vitamin A** has caused the homeotic HOX genes 1-4 to become expressed in groups of cells that usually do not express these genes. It is well known that high doses of vitamin A taken by women early during pregnancy can disturb the regulation of HOX genes even in the human embryo and cause severe malformations.



# Hierarchy of genes in *Drosophila* development



Development  
of the number  
of segments

Development  
of the features  
of the  
segments

# Hox genes in the Animal Kingdom

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Sponges



Sponges are the simplest animals, with bodies that are not organized along a body axis.

Anemones



Anemones have a primitive body axis, showing radial symmetry.

Flatworms



The other animals shown in this figure have a more complex form of symmetry called bilateral symmetry, meaning that their bodies are organized along a well-defined anteroposterior axis, with right and left sides that show a mirror symmetry. Such organisms are called bilaterians. Flatworms are very simple bilaterians.

Insects



Invertebrates such as insects are structurally more complex than flatworms, but less complex than organisms with a spinal cord.

Simple chordates



Animals with spinal cords are known as chordates. The simple chordates lack bony vertebrae that enclose the spinal cord.

Mammals



The vertebrates, such as mammals, have vertebrae and possess a very complex body structure.

