## ACUTE CHOLECYSTITIS

Supervisor: Dr. Abbas Mohammed

By: Abdullah Al-Rebdi, Hamad Bin Shaheen

- Anatomy
- Bile
- Stone formation & types
- Acute cholecystitis
  - Calculus
  - Acalculus
- Sign & symptoms
- Differential diagnosis
- Diagnostic investigation
- Complication
- Management
- Summery
- Questions

The gallbladder is located in the right upper quadrant of the abdomen beneath the liver.

The cystic duct exits at the neck of the gallbladder and joins the common hepatic duct to form the common bile duct.

CBD empty into the duodenum at the ampulla of Vater. This is surrounded by the sphincter of Oddi, which regulates bile flow into the duodenum .

#### Blood supply is from the cystic artery.

Celiac A.  $\longrightarrow$  Hepatic A.  $\longrightarrow$  Rt. Hepatic A.  $\longrightarrow$  Cystic A.

#### The cystic vein drain directly into portal vein.

#### Nerve Supply :

Sympathetic and parasympathetic vagal fibers the celiac plexus.

- Lymph Drainage:
- The lymph drains into a cystic lymph node situated near the neck of the gallbladder. From here, the lymph vessels pass to the hepatic nodes along the course of the hepatic artery and then to the celiac artery.







Bile produced in the liver is stored in the gallbladder.

#### The function of bile is emulsify FAT

Cholecystokinin stimulates gallbladder contraction and release of bile into the duodenum.

## Bile

The *spiral valves of Heister* in the cystic duct prevent bile reflux into the gallbladder.

Bile composed of

- Cholesterol
- Lecithin (phospholipid)
- Bile acid
- Bilirubin

Imbalance of cholesterol and its solubilizing agents, bile salts and lecithin concentrations

If hepatic cholesterol secretion is excessive then bile salts and lecithin are "overloaded", supersaturated cholesterol precipitates and can form gallstones Cholesterol stones

Pigment stones

Mixed stones

#### **Cholesterol Stones**



#### Pigment stones



#### Mixed stones:



# Acute Cholecystitis

## Inflammation of the gallbladder, resulting from :

- Obstruction of cystic duct by gallstone( 80%)
  )
- 2. Acalculous ( 20% )

Cholecystitis with stones most commonly blocking the cystic duct directly. This leads to inspissation (thickening) of bile, bile stasis, and secondary infection by gut organisms.

The most common organisms cultured during acute cholecystitis are Escherichia coli, Klebsiella, enterococci, Bacteroides fragilis, and Pseudomonas. The gallbladder shows congestion, thickening of the wall by edema and mucosal ulceration.

#### Thickened gallbladder wall

## Stone in cystic duct

#### Multifaceted stones in the gallbladder

网络哈拉克德的印度马克德尔拉拉马德的加克多纳尔兰拉克马克尔拉斯加利亚

#### Common bile duct -

## Pathogenesis:



Acute cholecystitis without gallstone may occur in a variety of condition it may be due to :

- Dehydration, prolonged fasting, TPN
- Systemic disease
- Generalized sepsis, trauma
- Kinking or fibrosis of the gallbladder
- Thrombosis of the cystic artery
- Sphincter spasm with obstruction of the biliary and pancreatic ducts
- Collagen vascular disease, DM, immunosuppressed

#### Hx

- □ Age 30-60 yrs
- $\square$  mostly  $\bigcirc$
- Sudden pain in RUQ, often radiate through the back to the tip of Rt. Scapula
- □ Pain is **continuous**, last >6 hrs
- Exacerbated by moving and breathing
- □ N/V
- Dark urine, pale stools, itchy skin in case of obstructive jaundice.

#### Hx:



right upper quadrant (with or without Murphy's sign and with or without a palpable mass)

(indicated by fever, concentrations of white blood cells and C-reactive protein above normal, and erythrocyte sedimentation rate above normal)

- Anorexia. Low grade fever ( < 38.5 C)
  - Tachycardia
  - Positive Murphy's sign
  - Palpable gallbladder (in 1/3 of patients)

## P/E

- Pt distressed by pain and lies quitely breathing shallowly.
- Tachycardia and Pyrexia
- □ Tenderness & guarding in Rt. Hypochondrium
- Boas' sign
- Murphy sign
- □ ± jaundice
- Rebound

#### Physical features of enlarged GB

- □ It appears from beneath the tip of the Rt. 9<sup>th</sup> rib
- Smooth & hemi-ovoid
- Moves with respiration
- There is no space between the lump and the edge of the liver
- Dull to percussion

- Perforating or penetrating peptic ulcer.
- Myocardial infraction
- Pancreatitis
- Hiatus hernia
- RLL pneumonia
- Appendicitis
- Hepatitis

- Leukocystosis
- Mild Hyperbillirubinemia
- $\Box$  Mild  $\uparrow$  AST, ALT, ALP
- Moderate ↑ Lipase & Amylase

## US

- Distended gallbladder
- Thickened gallbladdr wall
- Pericholecystic fluid collection
- □ US Murphy's sign ( + in 98% of pts)

Biliary scintigraphy (hydroxyiminodiacetic acid) (HIDA) scan:

Is the gold standard investigation when the diagnosis remains in doubt after ultrasound scanning.

## (HIDA) scan:

The patient is given an intravenous injection of radiolabelled hydroxyiminodiacetic acid and then the abdomen is scanned; in patients with acute cholecystitis, the gallbladder lumen will not take up any radioactive isotope one to two hours after injection and therefore the gall bladder will not be visible on the scan.

## (HIDA) scan:



- Cholecystentiric fistula formation
- Gallstone illeus
- Perforation (may cause localized abscess or generalized peritonitis [ after 3 days of onset ] )
- Empyema of gallbladder.
- Gangrene

## Mx

- □ NPO
- □ IVF
- IV Analgesia (pethidine)
- IV Abx (cefuroxime)
- □ NGT
- Parental Analgesics
- Optimal Rx is cholecystectomy within 3 days of the onset of symptoms.
- Pts who are poor candidates for surgery can undergo cholecystostomy (drainage of gallbladder contents)

## Calot's Triangle

- Superiorly: Lower edge of the liver
- Laterally: Cystic Duct
- Medially: Common Hepatic Duct

#### Acute Cholecystits vs. Billiary Colic

Duration, Symptoms ?

#### Surgery



FADAM.

#### Laproscopic



## Questions



- 1- Browse's Introduction to The Symptoms and Signs of Surgical Diseases, Fourth Edition.
- 2- Essentials of General Surgery, Lawrence, 4<sup>th</sup>
  Edition.
- 3- Surgical Recall