

ECP PSK



Enhanced External Counterpulsation (EECP)

*Goodbye to surgery
Go for Natural bypass with EECP.*

Chongqing PSK-Health Sci-Tech Development Co., LTD

Add: Room 14-8, NO. 5 Yanghe Sancun, Jiangbei

District, Chongqing, 400020, China.

Tel: 86-86833888

Fax: : 86-23-63834594

Email: export01@eecp.com.cn

Website: <http://www.eecp.com.cn>



Indications

- ✓ Stable & Unstable Angina
- ✓ Congestive Heart Failure
- ✓ Acute Myocardial infarction
- ✓ Cardiogenic Shock

TI model



T model





TM model



Paediatric Type for children


What is EECp ?

- EECp-Enhanced External Counterpulsation is an FDA-approved, non-invasive, non-surgical and outpatient medical therapy for the treatment of angina, congestive heart failure, acute myocardial infarction, and cardiogenic shock. During the treatment, blood pressure cuffs, wrapped around your legs, squeezed and released in sync with your heartbeat, promoting blood flow throughout your body and particularly to your heart. In the process, EECp develops new pathways around blocked arteries in the heart by expanding networks of tiny blood vessels ("collaterals") that help increase and normalize blood flow to the heart muscle. For this reason, it is often called the **NATURAL BYPASS**.

- Numerous clinical trials have shown EECp therapy to be safe and effective for patients with refractory angina with a clinical response rate averaging 70-80%, which is sustained up to five years.

What does EECF do?


EECP therapy consists of a system of three sets of inflation pressure cuffs wrapped around the calves, thighs and hips and then rapidly inflated and deflated in sync with the patient's heartbeat. Using a heart monitor, the cuffs sequentially inflate with air when the heart relaxes and deflate when the heart pumps. The timing of the inflation and deflation makes it easier for the heart to pump and increases blood supply to the heart. Studies show that 75% of patients treated with a single course of EECF experience a reduction in their angina and increased exercise tolerance.



EECP Principles of Operation


Diastolic Inflation

Sequentially inflate three sets of cuffs at the end of systole



Systolic Deflation

Simultaneously deflate all three sets of cuffs at the end of diastole



Diastolic Augmentation



Increase Coronary Perfusion

Increase Venous return



Increase Cardiac Output

Systolic Unloading





Reduce Cardiac Workload



Increase Cardiac Output

Myocardial Energy Demand and Supply



Time Tension Index

Workload of the heart is related to myocardial oxygen consumption



Diastolic Pressure Time Index

Energy supply to the myocardium in proportion to coronary perfusion pressure

Benefit of EECp

- Angina
- Angioplasty (PTCA) or Bypass surgery (CABG).
- CAD – Coronary Artery Disease
- CHD – Coronary Heart Disease
- PVD – Peripheral Vascular Disease
- ED – Erectile Dysfunction
- Cardiac chest pain
- Congestive heart failure
- Peripheral vascular disease
- Cardiomyopathy
- Peripheral neuropathy
- cerebral palsy
- Intestinal vascular insufficiency
- Edema, or venous insufficiency
- Chronic fatigue syndrome

Other benefits:

- Stroke
- kidney disease
- Parkinson's disease
- Memory disorders
- Diabetes and Diabetic Neuropathy
- High blood pressure
- Macular degeneration
- Other circulatory diseases
- hearing loss and tinnitus
- vision impairment
- autoimmune diseases (including Raynaud's phenomenon)
- rheumatic disease
- restless leg syndrome.
- Lymphatic System
- Sports Enhancement
- Anti-aging

CONTRAINDICATIONS

- Arrhythmias that interfere with machine triggering (need rate controlling)
- Bleeding diathesis (INR must be < 2.5)
- Active thrombophlebitis
- Severe lower extremity peripheral vascular disease
- Presence of a documented aortic aneurysm requiring surgical repair
- Pregnancy

PRECAUTIONS

- Patients with blood pressure higher than 180/110 mmHg should be controlled prior to treatment.
- Patients with a heart rate of more than 120 bpm should be controlled prior to treatment.
- Patients at high risk of complications from increased venous return should be carefully chosen and monitored during treatment. Decreasing cardiac afterload by optimizing cuff inflation and deflation timing may help minimize increased cardiac filling pressures and the possibility of pulmonary congestion due to increased venous return.
- Patients with clinically significant valvular disease should be carefully chosen and monitored during treatment with enhanced external counterpulsation. Certain valve conditions, such as significant aortic insufficiency or severe mitral or aortic stenosis, may prevent the patient from obtaining benefit from diastolic augmentation and reduced cardiac afterload in the presence of increased venous return.

EECP Treatment Regimen

■ Standard Treatment Time

- 5 daily 1 hour treatments per week over 7 weeks for a total of 35 hours or
- 2 x 1 hours daily over 3½ weeks for 35 hours total

■ Extension

- 7% from IEPR-2 had extended their 35 hours by 10.3 ± 9.8 hours because of persistent angina (67%), patient's preference (41%), physician's (40%)
- Extension is safe and patients continued to benefit with significant incremental improvement in symptoms and functional class

■ Repeat Therapy

- 18% of the patients having completed their initial course of 35 hours of EECP undergo retreatment within 2 years
- Common reasons for retreatment are recurrent angina, persistent angina
- About 13% of the patients failed to complete their initial 35 hours course of EECP because of patient's choice and adverse clinical events
- 30% of those who failed returned within 1 year for retreatment
- At retreatment, patients realized a benefit similar to patients who respond to a first course, with 70% improved by at least one CCS angina class, decreased angina episodes and nitroglycerin use.

F.A.Q.

■ How do I personally know EECP treatment has helped me?

- Patient can walk more distance without chest pain
- Patient would have fewer or no angina
- Episodes of angina would be less painful
- Patient can return to work and can participate in their active life style once again
- Patient would be more energetic and confidence.

■ Are there any risks or side effects of EECP?


EECP is safe. Occasionally, some patients experience mild skin irritation under the areas of the blood pressure cuffs. Experienced EECP therapists address this irritation by using extra padding to make the patient comfortable. Some patients experience a bit more fatigue at the beginning of their course of treatment, but it usually subsides after the first few sessions. In fact, patients typically feel energized by EECP.

■ How does EECP compare to angioplasty or bypass surgery?

- The five-year outcomes for EECP patients are virtually the same as for angioplasty and bypass surgery patients.

- **What are advantages of EECP?**
- EECP is non-invasive, simple, safe, risk-free and cost effective treatment without surgery or hospital stay. Patient can take this treatment without disturbing his/her daily routine life.
- **Is EECP possible after angioplasty or bypass surgery?**
- Yes, When the symptoms recurs or where the results of these procedures are inadequate or for additional benefit for a better and more active lifestyle.
- **Long term benefits**
- Data from the International EECP Patient Registry (IEPR) by the University of Pittsburgh's Graduate school of Public Health, USA suggest that the reduction in angina following EECP treatment is frequently sustained for up to 2 years post treatment. Patient follows up in many studies suggest that benefits of EECP persist for up to 5 years or more.








PRE EECp

POST EECp

EECP helps grow new collaterals for blood to flow, like a natural bypass around blocked arteries.



Very dark due to reduced blood flow
Extremely bright due to all of the new vessels feeding the heart


Clinical Evidence

The International EECF Patient Registry (IEPR) has provided data on over 5,000 patients demonstrating therapeutic outcomes and duration of benefit. Functional scores were graded using CCS angina score - classes I (mild) to IV (severe).

Summary:


- After 24 months follow up 31% of patients recorded being angina free compared to 0% at the start of the study.
- 82% of patients improved after EECF by one or more CCS class
- 43.9% of patients improved after EECF by two or more CCS classes
- Benefits were sustain over the 24 month follow up.

Improvement Maintained At 3-Year Follow Up




Visible evidence of hemodynamic effect on actual EECF patient

- Systolic unloading reduces energy requirements of the heart
- Dramatic diastolic augmentation (equal to or greater than intraaortic balloon pump)



Doppler echo of the descending aorta during EECF treatment

- Increased retrograde diastolic and enhanced systolic flow



Problems in Treating Heart Failure

- As the society aging, and the mortality rate from patients suffering from myocardial infarction decreases, the number of patients with heart failure will increase at a much faster pace, placing much more stress on the healthcare system
- Currently there is no effective therapy for heart failure
- The mortality rate for heart failure remain high (2001: 53,000 death/year)
- The difficulty in defining heart failure is because it is not a single organ disease but a systemic disease
- We require a treatment that not only improve the cardiac function but provide systemic pathophysiological benefits

Treatment of Heart Failure

Objectives:

- Symptomatic Improvement
- Preventing transition of asymptomatic cardiac dysfunction to symptomatic HF
- Preventing worsening of symptoms and/or functional limitations of HF
- Reducing ER visits and hospitalizations
- Reducing mortality


Pharmacological therapy:

- Prevention – hypertension, diabetes and underlying causes


Device:

- Enhanced External Counterpulsation (EECP)
- Pacemaker
- Implantable cardioverter defibrillator in pts with ventricular tachycardia or ventricular fibrillation
- Ventricular assist device / Artificial heart
- Ultrafiltration
- Cardiac Resynchronization therapy; pts with abnormal conduction


EECP Improves Each Major Pathophysiologic Feature of Heart Failure



Hemodynamics of the Heart



$$\text{Flow} = \frac{\text{Pressure}}{\text{Resistance}}$$



Energy balance
 Supply: Diastolic Pressure Time
 Demand: Systolic Pressure Time


The positive influence to blood circulation by EECp

Reduce systolic resistance




Increase cardiac output

Duplex echocardiography Descending Aorta



Lawson, Hui: J of Critical Illness 2000;5:629-636



- **EECP TREATMENT CAN ALSO BE USED FOR PREVENTIVE PURPOSES**
- Patients who had successful balloon-stent application and/or bypass operation because of significant narrowings in their coronary arteries and are asymptomatic but are candidate for early re-stenosis and occlusions due to having
- small coronary arteries (as mostly seen in diabetic and woman), should receive EECP therapy, in order to prolong the asymptomatic period provided by these interventions.
- Patients with coronary artery disease who have no symptoms but have mild or moderate narrowings in their coronary arteries, and also patients who have no documented coronary artery disease but have strong family history and many risk factors and therefore candidate for coronary artery disease, can get benefit from EECP's preventive effects.
- In summary, EECP is a useful treatment for every stage of coronary artery disease beginning from the presence of risk factors without documented disease to the advanced stage of the disease with severe symptoms refractory to other treatment modalities.

EECP

Safest, Non-surgical, FDA approved.



P-ECP/TI (All-in-One
Touch screen type)



P-ECP/TI (All-in-One Laptop
type)

Characteristics:

- Passed by CE certificate and Bio-compatibility test.
- It is based on Modern Medicine, Bio-Clinic Medicine, and Scientific Precise Data.
- It adopts the latest computer, modern control technique, and software control system.
- It adopts Germany Air Compressor and electromagnetic value only for EECp with proprietary intellectual property rights.
- It adopts scientific designed system to reduce noise and heat.
- It adopts integration of equipment (All in One) design.
- Its shape is based on the ergonomic design.

Transport and Storage Environment

- Temperature 14 to 104 (-10°C to 40°C)
- Relative Humidity $\leq 80\%$
- Atmospheric Pressure 0.1013MPa Atmospheric Free of corrosive gas

Dimension and Weight

- L*W*H 2150mm×840mm×600mm
- Net Weight 178 kg

Power Requirement

- AC Single Phase:220V±22V 50Hz/60Hz
- Maximum Power: 2.6KVA

Model: P-ECP/T (split type)



Characteristics:

- Design of air compressor and treatment bed separately, the air compressor comes with soundproof, place the air compressor and treatment bed in different rooms to make the patient completely free from noise impact during treatment;

Transport and Storage Environment

- Temperature 14 to 104 (-10°C to 40°C)
- Relative Humidity $\leq 80\%$
- Atmospheric Pressure 0.1013MPa
- Atmospheric Free of corrosive gas

Dimension and Weight

- Treatment Bed L×W×H 1980mm×800mm×700mm
- Net Weight 117kg
- Soundproof Box (Including Air compressor) L×W×H 720mm×600mm×730mm
- Net Weight 75kg
- Bedside Desk L×W×H 500mm×400mm×680mm
- Net Weight 18kg

Power Requirement

- AC Single Phase: 220V±22V 50Hz/60Hz
- Maximum Power: 2.6KVA

Model: P-ECP/TM (Movable type)



Characteristics:

- It is movable type with small space which make the treatment more easily.
- The air compressor adopted in the machine is Becker sliding vane rotary vacuum pump which is made in Germany and is full of gas, low noise and light.
- All the indexes of boxes of product have passed the CE Certificated and the test of bio-compatibility.
- It adopts the electromagnetic value with independent control, the characteristic of inflation and deflation are more excellent.
- It adopts special noise reduction and heat dissipation design.

Transport and Storage Environment

- Temperature 14 to 104 (-10°C to 40°C)
- Relative Humidity $\leq 80\%$
- Atmospheric Pressure 0.1013MPa
- Atmospheric Free of corrosive gas

Dimension and Weight

- L×W×H 780mm×515mm×960mm
- Net Weight 121kg

Power Requirement

- AC Single Phrase: 220V \pm 22V
50Hz/60Hz
- Maximum Power: 2.6KVA



Split Model: Pediatric Type for children



Characteristics:

- CE approval and passed biocompatibility tests;
- Based on modern medicine, bio-clinical medicine, high-precision scientific data;
- Professional design in accordance with the characteristics of children such as the size, appearance, comfort;
- Bladder design specifically for Children to prevent from all kinds of circuit malfunction;
- Scientific design of cooling, noise reduction system;
- Design of air compressor and treatment bed separately, the air compressor comes with soundproof, place the air compressor and treatment bed in different rooms to make the patient completely free from noise impact during treatment;

Treatment Bed

L×W×H: 1600mm×710mm×610mm

Soundproof Box (including air compressor)

L×W×H: 720mm×600mm×730mm

Net weight: 75kg

Bedside Desk

L×W×H: 500mm×400mm×680mm

Net weight: 18kg

Media reports about Enhanced External Counterpulsation

Bypassing the surgeon (不需要外科的“心脏搭桥”！)

MARCH 8, 2004

TIME

The Big Squeeze

Patients lie down during the procedure, which lasts an hour and is performed once a day, five times a week, for seven weeks. (The cost is about \$6,000, compared with as much as \$60,000 for bypass surgery.) The pneumatic cuffs are timed to inflate in progression—starting with the section around the calves—when the heart reaches its resting phase between beats. As each cuff inflates, it squeezes blood out of the legs and back to the heart. “It feels like a deep muscle massage,” says Dr. Debra Braverman, who administers ECEP to patients in Philadelphia. The most common side effect is chafing of the skin—usually prevented by wearing elastic clothing. Folks who have very high blood pressure, valve disease, phlebitis (inflammation of a vein) or are pregnant should not undergo ECEP.

Intriguingly, recent studies suggest that the heart responds to this extra flow of blood by producing tiny blood vessels to better nourish the heart. That may be why the benefits of ECEP often last several years. ECEP may also be useful in other hard-to-treat conditions, like congestive heart failure. “It’s probably underused,” says Dr. William Lawson, director of interventional cardiology at Stony Brook University Hospital in New York. That may change as the benefits of squeeze therapy become better known. ■

How an unlikely blood-pressure device relieves the chest pain of heart disease

By CHRISTINE GORMAN

IF YOU HAVE HEART DISEASE, you probably know all about statins and beta blockers, angioplasty and bypass surgery, and the benefits of regular exercise and a diet that’s low in saturated fat. But have you heard about strapping oversize blood-pressure cuffs to your legs and buttocks and pulsing them in synch with your heartbeat? The idea behind this wacky-sounding treatment, known as enhanced external counterpulsation (ECEP), is to decrease the demand on an ailing heart by helping it push blood through the body. But perhaps the oddest thing about ECEP is that it works amazingly well to relieve chest pain, or

angina, in many patients. First approved by the Food and Drug Administration in 1995, ECEP is most often used in folks with stable angina—the kind that often lasts five minutes or less, is brought on by physical exertion and is usually relieved by drugs like nitroglycerin. (Unstable angina tends to be severe, occurs suddenly or unexpectedly, often while a person is at rest, and requires immediate attention as it may be a sign of a heart attack.) Unfortunately, stable angina isn’t always easily controlled with medications, and some people just aren’t good candidates for angioplasty or surgery. That’s where ECEP comes in.

Cardiovascular News

Currents & Countercurrents

E. H. Squibb & Sons, Inc. has announced that it received FDA approval for a once-a-day diuretic for the management of hypertension. The combination tablet called Countipil consists of nifedipine (Corgard, Squibb) and furosemide (Lasix, Squibb) in a 100-mg/40-mg combination.

A survey of patients with confirmed acute intermittent porphyria (AIP) has revealed a significant association (p < 0.01) between this form of porphyria and chronic renal failure arising in nephrotic syndrome. Failure, p 4.

Some people are perhaps addicted to stress. They seek stressful situations because these raise their testosterone levels. The release of catecholamines, suggestions that this may be the root of some individuals.

Having shown that a steamed artichoke was the original perpetrator of the Pittman-Moreland scam by a man of the cloth, the historians of science have

While the skeletal morphology of North rhombomylitis or anorexia nervosa may result in significant elevations of serum creatinine, and still (continued on page 3)

Dr. Taro Takemi, a recent president of the Japan Medical Association, has proposed that the atomic bombs dropped over Hiroshima and Nagasaki may be said to have saved the Japanese from even more devastating carnage. While the two bombs killed, at least one million

A study aimed at analyzing the apparent greater longevity of centenarians did also revealed that cigarette smoking is the sole key to the difference. The study was done by Dr. Dean Gorman.

Sequenced Counterpulsation Described

CLEVELAND—Workers at the Cleveland Clinic are considering the feasibility of introducing a Chinese method of sequential external counterpulsation (SECP) in the treatment of acute MI and angina.



Patients are used in support of external counterpulsation technique over others that had an impact on health and safety.

Large Series Shows Snaked Bypass Grafting Indicated Even in Some Asymptomatic Patients

SAVRE, PENNSYLVANIA—Based on an analysis of 973 consecutive patients who underwent coronary bypass by the snaked graft technique, researchers here conclude that “surgery should be done on almost any asymptomatic, stable patient under the age of about 60 who is free of major organ disease (if he is within 50 pounds of normal weight and has an angiographic demonstration of coronary lesions which expose him to risk of myocardial infarction or death, whether he is symptomatic or not.”

Evidence Suggests Presumed Renal Damage From Captopril Therapy Dubious and Waning

CLEVELAND—The glomerular basement membrane deposits in capillary-mediated membranous glomerulonephritis apparently are not readily reversible and may be associated with persistent proteinuria, a recent study suggests. However, Dr. Stephen C. Textor announced in an interview with Cardiovascular News that “the late progressive renal failure seen in these patients is very likely to be related to pre-existing progressive arteriosclerosis/diabetes due to vascular disease itself.”

Marker May Foretell Ischemic Damage

ANN ARBOR, MICHIGAN—The leukocyte migration inhibition assay (LMI) is a relatively sensitive but nonspecific marker of ischemic myocardial infarction.

Dr. David K. Das and others at the University of Michigan Medical Center, Ann Arbor, evaluated the LMI assay for prognostic value in patients with acute myocardial infarction.

Buerger’s Disease I: Be Immune Disorder

TELEHASSETON, TEXAS—Patients with thromboangiitis obliterans (TAO) have been found to have higher cellular sensitivity to human T and T_H1 cytokines than healthy subjects or patients with atherosclerosis.

Depressed LV Function Among Effects of F

SCIENTIFARD, NEW YORK—Patient recovering from high fever may have a significant improvement in ventricular function, whether or not has coronary artery disease. C

By examining the left ventricular performance of high fever patients, researchers here have found that the fibrinolytic depression of function before the changes cause fibrinolytic.

During the course of ongoing treatment of 23 hypertensive patients, Dr. Textor and others at the Cleveland Clinic Foundation reported in the *American Journal of Medicine* (74:707, 1983) two cases of nephritic syndrome and membranous glomerulonephritis were encountered.

In both patients, “based on pathology,” Dr. Textor said, “the membranous changes, although persistent, were quite unrelated to their renal failure. The

Contents, page 3

Special Section: Advances in V See Page 4

BACK OF THE BOOK

MEDICINE

Bypassing the Surgeon

A new technique could simplify heart treatment



Suiting up: Pressure may succeed where surgery fails

Two years ago, Ann Clement started talking to funeral directors about her husband, Pierre. His kidneys had failed, and his diseased coronary arteries were rapidly starving his heart. At 61, the Long Island gift-shop owner was too weak to walk, too frail to survive surgery. Only morphine made his chest pain bearable. “His condition was zero,” Mrs. Clement recalls. “The doctors told us he was going to deteriorate and die.”

Today, Pierre Clement is off morphine, out of his wheelchair and walking two miles every morning. What changed his life was not a heroic operation or a costly new drug but a pair of pressurized pants. Clement is one of 18 U.S. heart patients pioneering a treatment called enhanced external counterpulsation. Rather than having his arteries catheterized or surgically bypassed, he spent 36 hour-long sessions strapped into a garment that works like a giant, pulsating blood-pressure cuff, painlessly forcing blood from the legs and hips into the chest. No one knows precisely how the treatment works or how long the effects last. But researchers at the Health Sciences Center in Stony Brook, N.Y., have recently shown that it can succeed where more costly and invasive treatments fail. If

larger trials confirm that finding, cardiology may never be the same.

Dr. Harry Sorooff, a heart surgeon at Stony Brook, conceived the counterpulsation technique 30 years ago at Harvard. During the 1960s and ‘70s, he and a collaborator designed various pulsating leg cuffs in attempts to force blood into the heart between beats. Unfortunately, their early devices lacked the power and precise timing the task demanded. Overshadowed by such flashy innovations as bypass surgery and balloon angioplasty (which uses a balloon-tipped catheter to reopen blocked arteries), counterpulsation fell into obscurity in the United States. In China, however,

proved by Thallium active dye to the heart, cent) regained returned technical mystery, t forcing blood ing or enlarging the vessels that p the disease. The researchers’ ling larger

Clinical research

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

American Heart Association® 
Learn and Live™

Left Ventricular Systolic Unload and Doppler Flow Du
Andrew D. Michaels, Mich
Circulation 2002;106:1237
DOI: 10.11
Circulation is published by the American Heart Association
Copyright © 2002 American Heart Association

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

American Heart Association® 
Learn and Live™

Enhanced External Counterpulsation Inhibits Intimal Hyperplasia by Modifying Shear Stress-Responsive Gene Expression in Hypercholesterolemic Pigs

Yan Zhang, Xiaohong He, Xiaolin Chen, Hong Ma, Donghong Liu, Jinyun Luo, Zhimin Du, Yafei Jin, Yan Xiong, Jianguo He, Dianqiu Fang, Kuijian Wang, William E. Lawson,

Cl

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION

American Heart Association® 
Learn and Live™

Enhanced External Counterpulsation Improves Peripheral Artery Flow-Mediated Dilation in Patients With Chronic Angina. A Randomized Sham-Controlled Study

Randy W. Braith, C. Richard Conti, Wilmer W. Nichols, Calvin Y. Choi, Matheen A. Khuddus, Darren T. Beck and Darren P. Casey

Circulation published online Oct 4, 2010;

DOI: 10.1161/CIRCULATIONAHA.109.923482

Technical Cooperation And Exchanges



EECP experts Prof. William Lawson



Prof Zheng zhensheng(middle, The father of EECP) and our CEOs



Association of cardiovascular club President Prof Fu dayi and our CEO



Our CEO in the First Academic Exchange Conference of EECP



JHui Ph.D John Hui and our CEO



Prof Cai Dawei(middle, the author of External Counterpulsation) and CEOs

EECP users




PSK EECF WORLDWIDE

In the foreign market, we exported to **more than 20 countries**. Established 3
branch offices: **India, Bangladesh, Thailand.**




EECP AROUND THE WORLD

1994 FDA(Food and Drug Administration)Certification 

1999 American Medicare 

2002 ACC/AHA Guideline
ACC(American College of Cardiology)
AHA(American Heart Association)  American Heart Association
Learn and Live 

2006 ESC Guideline
ESC (European Society of Cardiology) 

2006 CMA Guideline
CMA (Chinese Medical Association) 

American College of Cardiology (ACC) / American Heart Association (AHA) (2002), European Society of Cardiology (ESC) (2006) and Chinese Cardiovascular Society(2006) all put ECP therapy into the guidelines for treatment of angina pectoris and coronary heart disease.

EECP Association China (MCA)



The 2nd International EECP Symposium




Launch of the International EECP

The International EECP Society (IEECPS) was created in October 2013 as an association of physicians and clinicians involved in the study, research, application and provision of Enhanced External Counterpulsation (EECP) Therapy. Enhanced External Counterpulsation (EECP) Therapy, is an FDA-cleared, non-invasive, treatment for the symptoms of cardiovascular diseases stable ischemic heart disease, angina and congestive heart failure and other. Clinical studies in over 160 published medical and scientific journal articles on the safety and efficacy of EECP therapy have demonstrated that EECP therapy eliminates or significantly reduces symptoms while also improving the quality of life for these patients. Follow up studies have shown these initial benefits to be maintained for 3-5 years. EECP therapy is covered by Medicare and most third-party payers in the U.S. and many countries globally.

The mission of the IEECPS is to promote excellence in the noninvasive treatment of cardiovascular diseases through physician education, research, increased patient awareness, representation, and the advancement of quality patient care with EECP therapy.

The 3rd International EECF Symposium



Contact us at:

Chongqing PSK-Health Sci-Tech Development Co., Ltd

Tel:+86-23-86837032 Fax: +86-23-63834594

Email: export05@eep.com.cn

*Add: Room 14-8, CITIC Bank Bldg., NO. 5 Yanghe Sancun,
Jiangbei District, Chongqing, China.Headquarter*

www.eep.com.cn