



# Колебания Солнца и звезд и температурные волны в фотосфере

Ю. Д. Жутжда

ИЗМИРАН

## Intrinsic stellar oscillation amplitude and granulation power density measured in photometry

### Solar reference values and CoRoT response functions

E. Michel<sup>1</sup>, R. Samadi<sup>1</sup>, F. Baudin<sup>2</sup>, C. Barban<sup>1</sup>, T. Appourchaux<sup>2</sup>, and M. Auvergne<sup>1</sup>

## Appendix A: Visibility coefficients

The visibility factor associated with a mode observed in intensity by an observed pointing in the direction of the star rotation axis is given by the following relation (see Dziembowski 1977):

$$S_{\ell}(\lambda) = \frac{\int_0^1 d\mu \mu I_{\lambda}(\mu) Y_{\ell}^0(\mu)}{\int_0^1 d\mu \mu I_{\lambda}(\mu)} \quad (\text{A.1})$$

where  $\mu \equiv \cos(\theta)$ ,  $\theta$  is the angle with respect to the  $z$  axis,  $I_{\lambda}(\mu)$  is the monochromatic specific intensity used to take into account the limb-darkening,  $\lambda$  is the wavelength and  $Y_{\ell}^m$  is the spherical harmonic associated with a mode with a degree  $\ell$  and azimuthal order  $m$ . The spherical system of coordinates  $(r, \theta, \phi)$  is chosen

Letter to the Editor

**Spectral darkening functions of solar  $p$ -modes – an effective tool for helioseismology**Y.D. Zhugzhda<sup>1</sup>, J. Staude<sup>2</sup>, and G. Bartling<sup>2</sup>**Calculation of Spectral Darkening and Visibility Functions for Solar Oscillations**C. Nutto · M. Roth · Y. Zhugzhda · J. Bruls ·  
O. von der Lühе

$$I_\nu(\nu, \mu) = I_{0\nu}(\nu, \mu) + \delta I_\nu(\nu, \mu) = \int_0^\infty e^{-\tau_\nu/\mu} B_\nu(\tau_\nu) \frac{d\tau_\nu}{\mu} +$$

$$\int_0^\infty e^{-\tau_\nu/\mu} \left[ \frac{dB_\nu(\tau_\nu)}{d \ln T_0} \frac{\delta T}{T_0} + \frac{\delta \kappa_\nu}{\kappa_{0\nu}} (B_\nu(\tau_\nu) - I_\nu(\tau_\nu, \mu)) \right] \frac{d\tau_\nu}{\mu},$$

$$I_\nu(\tau_\nu, \mu) = \int_0^{\tau_\nu} e^{(\tau_\nu - \tau'_\nu)/\mu} B_\nu(\tau'_\nu) \frac{d\tau'_\nu}{\mu},$$

Функция  
потемнения

Флуктуации прозрачности

Адиабатическое приближение

$$\frac{\delta \kappa_\nu}{\kappa_{0\nu}} = \left( \frac{\partial \ln \kappa_{0\nu}}{\partial \ln T} \right)_\rho \frac{\delta T}{T_0} + \left( \frac{\partial \ln \kappa_{0\nu}}{\partial \ln \rho} \right)_T \frac{\delta \rho}{\rho_0}$$

$$\frac{\delta T}{T_0} = (\gamma - 1) \frac{\delta \rho}{\rho_0},$$

$$\mu = \cos \theta, \theta -$$

Полярный угол

$$\frac{\delta T}{T_0} = \text{const.}$$




Локальное приближение

После интегрирования по диску находим относительную функцию видимости


$$\frac{\delta F_\nu / F_{0\nu}}{\delta T / T_0} = \left\{ \int_0^\infty \left[ \frac{dB_\nu(\tau_\nu)}{d \ln T} E_2(\tau_\nu) - \left( \frac{\partial \ln \kappa_\nu}{\partial \ln T} \right)_\rho \int_{\tau_\nu}^\infty \frac{dB_\nu(\tau'_\nu)}{d\tau'_\nu} E_2(\tau'_\nu) d\tau'_\nu - \right. \right. \\ \left. \left. - \frac{1}{\Gamma_3(\tau_\nu) - 1} \left( \frac{\partial \ln \kappa_\nu}{\partial \ln \rho} \right)_T \int_{\tau_\nu}^\infty \frac{dB_\nu(\tau'_\nu)}{d\tau'_\nu} E_2(\tau'_\nu) d\tau'_\nu \right] d\tau_\nu \right\} \left[ \int_0^\infty B_\nu(\tau_\nu) E_2(\tau_\nu) d\tau_\nu \right]^{-1}$$

Функция видимости в локальном приближении может быть вычислена, так как она зависит только от параметров невозмущенной атмосферы


## Производная от прозрачности по температуре



## Производная от прозрачности по плотности





Сплошная линия – функция видимости, длинные штрихи – вклад от функции Планка, короткие штрихи и пунктир – вклады от флуктуаций прозрачности из-за флуктуаций температуры и плотности




# Обратная задача

$$\left\{ \int_{\nu_{n1}}^{\nu_{n2}} f_n(\nu) \int_0^{\infty} \frac{\delta T}{T_0} \left[ \frac{dB_\nu(\tau_\nu)}{d \ln T} E_2(\tau_\nu) - \left[ \left( \frac{\partial \ln \kappa_\nu}{\partial \ln T} \right)_\rho + \frac{1}{\Gamma_3(\tau_\nu) - 1} \left( \frac{\partial \ln \kappa_\nu}{\partial \ln \rho} \right)_T \right] - \int_{\tau_\nu}^{\infty} \frac{dB_\nu(\tau'_\nu)}{d \tau'_\nu} E_2(\tau'_\nu) d \tau'_\nu \right] d \tau_\nu d \nu \right\} \left[ \int_{\nu_{n1}}^{\nu_{n2}} f_n(\nu) \int_0^{\infty} B_\nu(\tau_\nu) E_2(\tau_\nu) d \tau_\nu d \nu \right]^{-1} = F_n^{(obs)},$$



Многоканальный фотометр ДИФОС

Кружки – ДИФОС, квадраты – SPM(SOHO)



THE ASTROPHYSICAL JOURNAL, 547:491-502, 2001 January 20  
 © 2001. The American Astronomical Society. All rights reserved. Printed in U.S.A.




LAGRANGIAN AND EULERIAN STRATIFICATIONS OF ACOUSTIC OSCILLATIONS THROUGH THE SOLAR PHOTOSPHERE

INÉS RODRÍGUEZ HIDALGO, BASILIO RUIZ COBO, MANUEL COLLADOS, AND LUIS R. BELLOT RUBIO

APPLICATION OF INVERSE METHODS FOR INVESTIGATION OF SOLAR BRIGHTNESS OSCILLATIONS

M. Stodilka

The Ivan Franko National University of Lviv,  
 Astronomical Observatory  
 8 Kyrylo and Mephodij Str., Lviv, Ukraine, 79005



# Функции потемнения




Рис. 6: Функции потемнения для р-мод с  $l = 0$  для четырех каналов фотометра ДИФОС с длинами волн  $\lambda = 357.6(a)$ ,  $494.5(b)$ ,  $630.7(c)$ ,  $866.3(d)$  нм. Номера кривых на графиках соответствуют номерам моделей с фиксированной оптической толщиной на графиках рис.5.



Y. D. ZHUGZDHA

Department of Cosmic Electrodynamics, IZMIRAN, Troitsk, Moscow Region, 142092, U.S.S.R.

*Solar Physics* **124**: 205–209, 1989.

© 1989 Kluwer Academic Publishers. Printed in Belgium.


YU. D. ZHUGZDHA

Laboratory of Cosmic Electrodynamics, IZMIRAN, Troitsk, Moscow Region, U.S.S.R.

*Astrophysics and Space Science* **95** (1983) 255–275. 0004-640X/83/0952-0255\$03.15.

© 1983 by D. Reidel Publishing Co., Dordrecht and Boston

# Температурные волны



Квази-адиабатическое приближение

$$k_a = \frac{\omega}{c_T} + i \frac{c_T}{2\chi c_s^2} (c_s^2 - c_T^2) \quad (\text{acoustic wave}),$$

$$k_T = (1 + i) \sqrt{\frac{\omega}{2\gamma\chi}} \quad (\text{temperature wave}),$$

$$|k_T| \ll |k_a|$$


$$k^4 - k^2 \left( \frac{\omega^2}{c_T^2} + \frac{i\omega}{\chi} \right) + \frac{i\omega^3}{\chi c_s^2} = 0$$

Квази-изотермическое приближение

$$k_a = \frac{\omega}{c_s} + i \frac{\omega^2 \chi}{2c_s} \left( \frac{1}{c_T^2} - \frac{1}{c_s^2} \right) \quad (\text{acoustic wave}),$$

$$k_T^2 = \frac{i\omega}{\chi} + \omega^2 \left( \frac{1}{c_T^2} - \frac{1}{c_s^2} \right) \quad (\text{temperature wave}),$$


$$|k_T| \gg |k_a|$$



diffuse approximation is not correct in this layer. The temperature conductivity of the solar photosphere is about  $10^{11} \text{ cm}^2 \text{ s}^{-1}$  and the wavelength of temperature waves is around

$$\lambda(\text{km}) = 10 \sqrt{P} \text{ (s)}, \tag{10}$$


where  $P$  is the period of a wave.



Five-minute oscillations in the solar photosphere must generate temperature waves due to a linear interaction. In the layer of interaction, the temperature waves are not damped over short distances because of reinforcing by acoustic waves. In the photosphere the temperature disturbances of low-frequency (Equation (8)) temperature waves are connected with the velocity amplitude by

$$\left| \frac{\delta T}{T} \right| \simeq (P \text{ (s)})^{3/2} \frac{V}{c_s} \tag{11}$$

The temperature fluctuations of the five-minute temperature waves are greater by a factor  $10^3$  than temperature fluctuations generated by pure acoustic  $p$ -modes of the same velocity amplitude.



# Взаимодействие р-мод и температурных волн

$$i\omega\rho' + w\frac{\rho_0}{H} + \rho_0u = 0,$$

$$i\omega\rho_0w + \frac{dp'}{dz} - \rho'g = 0,$$

$$i\omega T' + (\gamma - 1)T_0u = \frac{Q'}{\rho_0c_v},$$

$$\frac{p_0}{T_0}T' = p' - \frac{p_0}{\rho_0}\rho';$$

Оптически толстая  
атмосфера

$$Q' = \lambda\left(\frac{d^2T'}{dz^2} - k_{\perp}^2T'\right) + \frac{d\lambda}{dz}\frac{dT'}{dz}.$$

$$\lambda = \lambda_0 \exp\left[\frac{(1-s)z}{H}\right].$$

Оптически тонкая  
атмосфера

$$Q' = -\rho_0c_vqT',$$

Изотермическая  
атмосфера  
Н-шкала высот

$$\frac{d^2}{dz^2}\left(\frac{Q'}{p_0}\right) + \frac{1}{H}\frac{d}{dz}\left(\frac{Q'}{p_0}\right) + \left(\frac{\omega^2}{c_*^2} - k_{\perp}^2\right)\frac{Q'}{p_0} - \frac{i\gamma\omega}{(\gamma-1)T_0} \times$$

$$\times \left[\frac{d^2T'}{dz^2} + \frac{1}{H}\frac{dT'}{dz} + \left(\frac{\omega^2}{c_0^2} + \frac{(\gamma-1)g^2k_{\perp}^2}{\omega^2c_0^2} - k_{\perp}^2\right)T'\right] = 0,$$

Уравнение имеет точное аналитическое решение в виде обобщенных гипергеометрических функций

