

Далее, из  $\text{rot } \vec{u} = \vec{0} \Leftrightarrow \vec{u} = \text{grad} f$ .  $\text{grad}(f + C) = \text{grad} f$ .

$$\vec{u} = \{P; Q, R\}, \text{rot } \vec{u} = \vec{0}, \text{ т.е. } \frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y} \in G \subset E_3$$

$$(x_0, y_0, z_0) \in G. \rightarrow f(x, y, z) = \int_{x_0}^x P(\xi, y, z) d\xi + \varphi(y, z), \text{ где, из } \frac{\partial f}{\partial x} = P(x, y, z)$$

$$Q(x, y, z) = \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} \int_{x_0}^x P(\xi, y, z) d\xi + \frac{\partial \varphi}{\partial y} = \left( \text{об-ва упр-лов,} \right) = \int_{x_0}^x \frac{\partial P(\xi, y, z)}{\partial y} d\xi + \frac{\partial \varphi}{\partial y} =$$

$$= \int_{x_0}^x \frac{\partial Q(\xi, y, z)}{\partial \xi} d\xi + \frac{\partial \varphi}{\partial y} = Q(x, y, z) - Q(x_0, y, z) + \frac{\partial \varphi}{\partial y}, \text{ т.е. } \frac{\partial \varphi}{\partial y} = Q(x_0, y, z) \Rightarrow$$

$$\Rightarrow f(x, y, z) = \int_{x_0}^x P(\xi, y, z) d\xi + \int_{y_0}^y Q(x_0, \eta, z) d\eta + \psi(z). \quad R(x, y, z) = \frac{\partial f}{\partial z} =$$

$$= \int_{x_0}^x \frac{\partial P(\xi, y, z)}{\partial z} d\xi + \int_{y_0}^y \frac{\partial Q(x_0, \eta, z)}{\partial z} d\eta + \psi'(z) = \int_{x_0}^x \frac{\partial R(\xi, y, z)}{\partial \xi} d\xi + \int_{y_0}^y \frac{\partial R(x_0, \eta, z)}{\partial \eta} d\eta + \psi'(z) =$$

$$= R(x, y, z) - R(x_0, y, z) + R(x_0, y, z) - R(x_0, y_0, z) + \psi'(z), \text{ т.е. } \psi'(z) = R(x_0, y_0, z) \Rightarrow$$

$$f(x, y, z) = \int_{x_0}^x P(\xi, y, z) d\xi + \int_{y_0}^y Q(x_0, \eta, z) d\eta + \int_{z_0}^z R(x_0, y_0, \xi) d\xi + C$$



$$f(x, y, z) = \int_{x_0}^x P(\xi, y, z) d\xi + \int_{y_0}^y Q(x_0, \eta, z) d\eta + \int_{z_0}^z R(x_0, y_0, \zeta) d\zeta + C$$

Пример.  $\vec{u} = -\frac{\vec{r}}{r^3} = \left\{ -\frac{x}{(x^2+y^2+z^2)^{3/2}}, -\frac{y}{(x^2+y^2+z^2)^{3/2}}, -\frac{z}{(x^2+y^2+z^2)^{3/2}} \right\}$   $\left( \begin{array}{l} \vec{r} = \{x, y, z\} \\ r = |\vec{r}| = \sqrt{x^2+y^2+z^2} \end{array} \right)$

$\text{rot } \vec{u} = \vec{0}$  (conserv.),  $\vec{u} = \text{grad } f$ ,  $f(x, y, z) = -\int_{x_0}^x \frac{\xi d\xi}{(\xi^2+y^2+z^2)^{3/2}} - \int_{y_0}^y \frac{\eta d\eta}{(x_0^2+\eta^2+z^2)^{3/2}} -$

$$- \int_{z_0}^z \frac{\zeta d\zeta}{(x_0^2+y_0^2+\zeta^2)^{3/2}} + C = \frac{1}{\sqrt{\xi^2+y^2+z^2}} \Big|_{\xi=x_0}^{\xi=x} + \frac{1}{\sqrt{x_0^2+\eta^2+z^2}} \Big|_{\eta=y_0}^{\eta=y} + \frac{1}{\sqrt{x_0^2+y_0^2+\zeta^2}} \Big|_{\zeta=z_0}^{\zeta=z} + C =$$

$$= \frac{1}{\sqrt{x^2+y^2+z^2}} - \frac{1}{\sqrt{x_0^2+y^2+z^2}} + \frac{1}{\sqrt{x_0^2+y^2+z^2}} - \frac{1}{\sqrt{x_0^2+y_0^2+z^2}} + \frac{1}{\sqrt{x_0^2+y_0^2+z^2}} - \frac{1}{\sqrt{x_0^2+y_0^2+z_0^2}} + C$$

$$= \frac{1}{r} - \frac{1}{r_0} + C = \frac{1}{r} + \text{const}$$

## О векторном потенциале

Потенциальное поле:  $\oint_{\Gamma} (\vec{u}, d\vec{r}) = 0$      $\operatorname{rot} \vec{u} = \vec{0}$      $\vec{u} = \operatorname{grad} f$

Соленоидальное поле:  $\oint_S (\vec{u}, d\vec{S}) = 0$      $\operatorname{div} \vec{u} = 0$     ?



## О векторном потенциале

Потенциальное поле:  $\oint_{\Gamma} (\vec{u}, d\vec{r}) = 0$      $\text{rot } \vec{u} = \vec{0}$      $\vec{u} = \text{grad } f$

Соленоидальное поле:  $\oint_S (\vec{u}, d\vec{S}) = 0$      $\text{div } \vec{u} = 0$      $\vec{u} = \text{rot } \vec{v}$

$\vec{v}$  - векторный потенциал

$$\vec{u} = \{P, Q, R\}, \quad \vec{v} = \{P_1, Q_1, R_1\}, \quad \text{rot } \vec{v} = \left\{ \underbrace{\frac{\partial R_1}{\partial y} - \frac{\partial Q_1}{\partial z}}_P, \underbrace{\frac{\partial P_1}{\partial z} - \frac{\partial R_1}{\partial x}}_Q, \underbrace{\frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y}}_R \right\} = \vec{u}$$

$$0 = \text{div } \vec{u} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} = \frac{\partial^2 R_1}{\partial x \partial y} - \frac{\partial^2 Q_1}{\partial x \partial z} + \frac{\partial^2 P_1}{\partial y \partial z} + \frac{\partial^2 R_1}{\partial x \partial y} + \frac{\partial^2 Q_1}{\partial x \partial z} - \frac{\partial^2 P_1}{\partial y \partial z}$$

$$P = \frac{\partial R_1}{\partial y} - \frac{\partial Q_1}{\partial z}, \quad Q = \frac{\partial P_1}{\partial z} - \frac{\partial R_1}{\partial x}, \quad R = \frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y} \quad \text{Положим } R_1(x, y, z) \equiv 0$$

$$\Rightarrow P = -\frac{\partial Q_1}{\partial z}, \quad Q = \frac{\partial P_1}{\partial z}, \quad R = \frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y}. \quad (x_0, y_0, z_0) \in G \Rightarrow P_1(x, y, z) \stackrel{z}{=} \int_{z_0}^z Q(x, y, \xi) d\xi$$

$$Q_1(x, y, z) = -\int_{z_0}^z P(x, y, \xi) d\xi + \varphi(x, y) \Rightarrow R(x, y, z) = \frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y} = -\int_{z_0}^z \frac{\partial P(x, y, \xi)}{\partial x} d\xi +$$

$$+ \frac{\partial \varphi}{\partial x} - \int_{z_0}^z \frac{\partial Q(x, y, \xi)}{\partial y} d\xi = \frac{\partial \varphi}{\partial x} - \int_{z_0}^z \left( \frac{\partial P(x, y, \xi)}{\partial x} + \frac{\partial Q(x, y, \xi)}{\partial y} \right) d\xi = \frac{\partial \varphi}{\partial x} + \int_{z_0}^z \frac{\partial R(x, y, \xi)}{\partial \xi} d\xi =$$

$$= \frac{\partial \varphi}{\partial x} + R(x, y, z) - R(x, y, z_0), \quad \text{т.е. } \frac{\partial \varphi}{\partial x} = R(x, y, z_0) \Rightarrow \varphi(x, y) = \int_{x_0}^x R(\xi, y, z_0) d\xi + \psi(y)$$

$\text{дерив} \equiv 0$

$$\text{Итак, } P_1(x, y, z) = \int_{z_0}^z Q(x, y, \xi) d\xi; \quad Q_1(x, y, z) = -\int_{z_0}^z P(x, y, \xi) d\xi + \int_{x_0}^x R(\xi, y, z_0) d\xi; \quad R_1(x, y, z) \equiv 0$$



Итак,  $P_1(x, y, z) = \int_{z_0}^z Q(x, y, \xi) d\xi$ ;  $Q_1(x, y, z) = -\int_{z_0}^z P(x, y, \xi) d\xi + \int_{x_0}^x R(\xi, y, z_0) d\xi$ ;  $R_1(x, y, z) \equiv 0$

Пример  $\vec{u} = \{y, z, x\}$   $\text{div } \vec{u} = 0$  Найдем  $\vec{v}$ :  $\vec{u} = \text{rot } \vec{v}$   $x_0 = y_0 = z_0 = 0$

$P_1(x, y, z) = \int_0^z \xi d\xi = \frac{z^2}{2}$ ;  $Q_1(x, y, z) = -\int_0^z y d\xi + \int_0^x \xi d\xi = \frac{x^2}{2} - yz$ ;  $R_1(x, y, z) \equiv 0$

$\vec{v} = \left\{ \frac{z^2}{2}, \frac{x^2}{2} - yz, 0 \right\}$   $\text{rot } \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{z^2}{2} & \frac{x^2}{2} - yz & 0 \end{vmatrix} = \{y, z, x\} = \vec{u}$ , т.е.  $\vec{v}$  - векторный потенциал



Итак,  $P_1(x, y, z) = \int_{z_0}^z Q(x, y, \xi) d\xi$ ;  $Q_1(x, y, z) = -\int_{z_0}^z P(x, y, \xi) d\xi + \int_{x_0}^x R(\xi, y, z_0) d\xi$ ;  $R_1(x, y, z) \equiv 0$

Пример  $\vec{u} = \{y, z, x\}$   $\operatorname{div} \vec{u} = 0$  Найдем  $\vec{v}$ :  $\vec{u} = \operatorname{rot} \vec{v}$   $x_0 = y_0 = z_0 = 0$

$P_1(x, y, z) = \int_0^z \xi d\xi = \frac{z^2}{2}$ ;  $Q_1(x, y, z) = -\int_0^z y d\xi + \int_0^x \xi d\xi = \frac{x^2}{2} - yz$ ;  $R_1(x, y, z) \equiv 0$

$\vec{v} = \left\{ \frac{z^2}{2}; \frac{x^2}{2} - yz; 0 \right\}$   $\operatorname{rot} \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{z^2}{2} & \frac{x^2}{2} - yz & 0 \end{vmatrix} = \{y, z, x\} = \vec{u}$ , т.е.  $\vec{v}$  - векторный потенциал

Задача. Проверить, что для поля  $\vec{u} = -\frac{\vec{r}}{r^3}$   $\operatorname{div} \vec{u} = 0$  и найти векторный потенциал



# Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|                                 |      |     |     |
|---------------------------------|------|-----|-----|
| $\nabla^{(2)}$ / $\nabla^{(1)}$ | grad | div | rot |
| grad                            |      |     |     |
| div                             |      |     |     |
| rot                             |      |     |     |



# Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

| $\nabla_{(2)}$ \ $\nabla_{(1)}$ | grad | div | rot |
|---------------------------------|------|-----|-----|
| grad                            | X    |     |     |
| div                             |      | X   | X   |
| rot                             | X    |     |     |



## Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}; \frac{\partial}{\partial y}; \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|   |      |          |     |
|---|------|----------|-----|
| $\nabla^{(2)}$ \diagdown $\nabla^{(1)}$ | grad | div      | rot |
| grad                                    | X    | $\Delta$ |     |
| div                                     |      | X        | X   |
| rot                                     | X    |          |     |

$$\operatorname{div} \operatorname{grad} f = \operatorname{div} \left\{ \frac{\partial f}{\partial x}; \frac{\partial f}{\partial y}; \frac{\partial f}{\partial z} \right\} = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \Delta f \quad (\text{однаковое})$$

$$\Delta - \text{оператор Лапласа, } \Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

скал. пр-ие

$$\text{Формально: } \operatorname{div} \operatorname{grad} f = (\nabla, \nabla f) = (\nabla, \nabla) f = \Delta f, \text{ т.к. } (\nabla, \nabla) = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

$\Delta$  можно применить и к векторному полю:  $\Delta \vec{u} = \Delta \{P; Q; R\} \equiv \{\Delta P; \Delta Q; \Delta R\}$



## Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|   |      |          |           |
|---|------|----------|-----------|
| $\nabla_{(2)}$ \diagdown $\nabla_{(1)}$ | grad | div      | rot       |
| grad                                    | X    | $\Delta$ | $\vec{0}$ |
| div                                     | X    | X        | X         |
| rot                                     | X    | X        | X         |

$$\text{rot grad } f = \begin{matrix} \vec{i} & \vec{j} & \vec{k} \\ \left\{ \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\} & \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} & \frac{\partial f}{\partial z} \\ \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} & \frac{\partial f}{\partial z} \end{vmatrix} & \left\{ \frac{\partial^2 f}{\partial y \partial z} - \frac{\partial^2 f}{\partial z \partial y}, \frac{\partial^2 f}{\partial x \partial z} - \frac{\partial^2 f}{\partial z \partial x}, \frac{\partial^2 f}{\partial x \partial y} - \frac{\partial^2 f}{\partial y \partial x} \right\} \\ = \text{rot} \left\{ \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\} & = & \left\{ \frac{\partial^2 f}{\partial x \partial x} - \frac{\partial^2 f}{\partial x \partial x}, \frac{\partial^2 f}{\partial y \partial y} - \frac{\partial^2 f}{\partial y \partial y}, \frac{\partial^2 f}{\partial z \partial z} - \frac{\partial^2 f}{\partial z \partial z} \right\} \\ & = & \{ 0, 0, 0 \} = \vec{0} \end{matrix}$$

Формально:  $\text{rot grad } f = [\nabla, \nabla f] = [\nabla, \nabla] f = \vec{0}$ , так  $[\nabla, \nabla] = \vec{0}$



## Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|                             |      |          |                |
|-----------------------------|------|----------|----------------|
| $\nabla_{(2)}$ \nabla_{(1)} | grad | div      | rot            |
| grad                        |      | $\Delta$ | $\vec{\sigma}$ |
| div                         |      |          |                |
| rot                         |      | 0        |                |

$$\operatorname{div} \operatorname{rot} \vec{u} = \operatorname{div} \operatorname{rot} \{P, Q, R\} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \operatorname{div} \left\{ \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}, \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}, \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right\} =$$






$$= \frac{\partial^2 R}{\partial x \partial y} - \frac{\partial^2 Q}{\partial x \partial z} + \frac{\partial^2 P}{\partial y \partial z} - \frac{\partial^2 R}{\partial x \partial y} + \frac{\partial^2 Q}{\partial x \partial z} - \frac{\partial^2 P}{\partial y \partial z} = 0$$

Формально:  $\operatorname{div} \operatorname{rot} \vec{u} = (\nabla, [\nabla, \vec{u}]) = 0$  (смешанное произведение)



## Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|                             |   |  |   |
|-----------------------------|---|--|---|
| $\nabla_{(2)}$ \nabla_{(1)} | grad  | div  | rot   |
| grad                        |  | $\Delta$   | $\vec{\sigma}$  |
| div                         | grad div  |  |  |
| rot                         |  | 0  |  |





$$\begin{aligned} \text{grad div } \vec{u} &= \text{grad div} \{P; Q; R\} = \text{grad} \left( \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) = \\ &= \left\{ \frac{\partial^2 P}{\partial x^2} + \frac{\partial^2 Q}{\partial x \partial y} + \frac{\partial^2 R}{\partial x \partial z}, \frac{\partial^2 P}{\partial x \partial y} + \frac{\partial^2 Q}{\partial y^2} + \frac{\partial^2 R}{\partial y \partial z}, \frac{\partial^2 P}{\partial x \partial z} + \frac{\partial^2 Q}{\partial y \partial z} + \frac{\partial^2 R}{\partial z^2} \right\} \end{aligned}$$

Формально:  $\text{grad div } \vec{u} = \nabla(\nabla, \vec{u})$  - далее не преобразуется



## Дифференциальные операции второго порядка

$f, \vec{u} \in C_2$   $\nabla = \left\{ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\}$  - оператор Гамильтона

|  |  |  |   |
|--|--|--|---|
| $\nabla_{(2)} \backslash \nabla_{(1)}$ | grad   | div  | rot   |
| grad                                   |   | $\Delta$   | $\vec{0}$   |
| div                                    | grad div   |  |  |
| rot                                    |  | 0  | grad div - $\Delta$   |

$$\text{rot rot } \vec{u} = [\nabla, [\nabla, \vec{u}]] = \nabla(\nabla, \vec{u}) - (\nabla, \nabla)\vec{u} = \text{grad div } \vec{u} - \Delta \vec{u}$$

Задача. Получить эту формулу неформально (т.е. выкладкой)