Западно-Казахстанский государственный медицинский университет имени Марата Оспанова

Кафедра: Доказательной медицины
Специальность: Акушерство и Гинекология

Тема : “Преждевременный дородовый разрыв плодных оболочек”

Выполнили : Кабдыгалиева А.
Хасенова С.
Группа : 105
Проверила : Назарбаева Р.К.

Актобе , 2019 год

- Клинический протокол преждевременный разрыв плодных оболочек. Дородовый разрыв плодных оболочек .
- Утверждено на Экспертной комиссии по вопросам развития здравоохранения Министерства здравоохранения Республики Казахстан протокол № 10 от «04» июля 2014 года

- **Ссылка :** https://diseases.medelement.com/disease/преждевременный-разрыв- плодных-оболочек/13991
- **Список разработчиков протокола:**
- 1) Укыбасова Талшын Мухадесовна – д.м.н., АО «ННЦМД», профессор, руководитель отдела акушерства и гинекологии.
- 2) Бапаева Гаури Биллахановна – д.м.н., АО «ННЦМД», доцент, начальник отдела науки и менеджмента научных исследований .
- 3) Иманкулова Балкенже Жаркемовна – к.м.н., АО «ННЦМД», врач акушер-гинеколог высшей категории отделения женских болезней.
- **Рецензент:**
  Сейдуллаева Лейла Алтынбековна – к.м.н., доцент кафедры интернатуры акушерства и гинекологии Медицинского Университета Астаны.

- **Недостатки протокола:**

1. В данном протоколе описывается , что в сроке 35 – 37 недель показана выжидательная тактика . Было бы лучше , если отказаться от выжидательной тактики в данном сроке , так как можно предотвратить осложнения оказывающие влияние и для матери и для ребенка , самое главное - **снизить частоту осложнении! ! !**

2.Отсутствие сносок, ссылок для быстрого перехода на использованную литературу.

3. В создании протокола участвовали только врачи акушер-гинекологи, при отсутствии их коллег: анестезиологов-реаниматологов.

4. Данный протокол №10 МЗ РК от 04.07.2014 требует пересмотра, в связи с истечением срока.

- **Р** – беременные с дородовым излитием околоплодных вод
- **I** –активная тактика
- **С** - выжидательная тактика
- **О** – снижение частоты осложнении (разрыв).

Вопрос:

Снижает ли частоту осложнении применение активной тактики по сравнению с отсутствием применения выжидательной тактики у беременных с преждевременным дородовым излитием околоплодных вод?

№ 10 МЗ РК от 04 июля 2014 года
Преждевременный разрыв плодных оболочек

pubmed - ничего не на | хориоамнионит - Publ | Google Переводчик | преждевременный ра | Преждевременный | ДМ Акушер-Гинеколог | ДЗ ДМ на 10.01.19г..pp

https://diseases.medelement.com/disease/преждевременный-разрыв-плодных-оболочек/13991

Войти

О проекте | Электронная клиника | MedElement | Для пациентов | Справочники

Преждевременный разрыв плодных оболочек

Также: Дородовое излитие околоплодных вод

РЦРЗ (Республиканский центр развития здравоохранения МЗ РК)
Версия: Клинические протоколы МЗ РК - 2014

Категории МКБ: Преждевременный разрыв плодных оболочек (O42)

Разделы медицины: Акушерство и гинекология

Общая информация

Версия для печати
Скачать или отправить файл

Краткое описание

Утверждено на Экспертной комиссии
по вопросам развития здравоохранения
Министерства здравоохранения Республики Казахстан
протокол № 10 от «04» июля 2014 года

Дородовый разрыв плодных оболочек (ДРПО) – спонтанный разрыв амниотических оболочек до начала регулярных сокращений матки в сроке 37 недель и более.
Преждевременный дородовый разрыв плодных оболочек (ПДРПО) – спонтанный разрыв амниотических оболочек до начала регулярных сокращений матки в сроке 22 –

Разделы | На главную

смертности связаны с ПДРПО: недоношенность, сепсис и гипоплазия легких [1,2]. Риск для матери связан, прежде всего, с

RU 22:34 09.01.2019

Шаг 2. Поиск информации.

Ключевые слова в Pubmed: prenatal rupture of membranes , Induction of prelabour , pregnant women , chorioamnionitis

pubmed - ничего не найдено. | prenatal rupture of membra | ДМ Акушер-Гинекологи 2019г.

https://www.ncbi.nlm.nih.gov/pubmed

PubMed.gov
US National Library of Medicine
National Institutes of Health

PubMed | prenatal rupture of membranes , Induction of prelabour , pregnant women , chorioamnionitis | Search

Create RSS Create alert Advanced Help

Article types clear
- ✓ Clinical Trial
- Review
- Customize ...

Text availability
- Abstract
- Free full text
- Full text

Publication dates
- 5 years
- 10 years
- Custom range...

Species clear
- ✓ Humans
- Other Animals

Clear all

Show additional filters

Format: Summary Sort by: Most Recent Send to

Best matches for prenatal rupture of membranes chorioamnionitis:

Mid-trimester preterm premature **rupture** of **membranes** (PPROM): etiology, diagnosis, classification, international recommendations of treatment options and outcome.
Tchirikov M et al. J Perinat Med. (2018)

Chorioamnionitis following preterm premature **rupture** of **membranes** and fetal heart rate variability.
Vandenbroucke L et al. PLoS One. (2017)

Tocolysis after preterm premature **rupture** of **membranes** and neonatal outcome: a propensity-score analysis.
Lorthe E et al. Am J Obstet Gynecol. (2017)

Switch to our new best match sort order

Search results

Items: 5

Filters activated: Clinical Trial, Humans. Clear all to show 138 items.

1. Single versus weekly courses of antenatal corticosteroids in preterm premature **rupture** of **membranes**.
Lee MJ, Davies J, Guinn D, Sullivan L, Atkinson MW, McGregor S, Parilla BV, Hanlon-Lundberg K, Simpson L, Stone J, Wing D, Ogasawara K, Muraskas J.
Obstet Gynecol. 2004 Feb;103(2):274-81.
PMID: 14754695
Similar articles

Filters: Manage Filters

Sort by: Best match | Most recent

Find related data
Database: Select
Find items

Search details
(prenatal[All Fields] AND ("rupture"[MeSH Terms] OR "rupture"[All Fields]) AND ("membranes"[MeSH Terms] OR "membranes"[All Fields]) AND ("chorioamnionitis"[MeSH Terms]

Search See more...

Recent Activity
Turn Off Clear
- prenatal rupture of membranes chorioamnionitis AND (Clinical Tria... (PubMed
- prenatal rupture of membranes chorioamnionitis AND (Clinical Tria... (PubMed

RU 23:17 09.01.2019

https://www.ncbi.nlm.nih.gov/pubmed/17617892

pubmed - ничего не на | chorioamnionitis - P | Google Переводчик | преждевременный ра | Преждевременный ра | ДМ Акушер-Гинеколо | ДЗ ДМ на 10.01.19г..pp

https://www.ncbi.nlm.nih.gov/pubmed

Similar articles

14. Correlation between initial neonatal and early childhood outcomes following preterm birth.
Manuck TA, Sheng X, Yoder BA, Varner MW.
Am J Obstet Gynecol. 2014 May;210(5):426.e1-9. doi: 10.1016/j.ajog.2014.01.046.
PMID: 24793722 Free PMC Article
Similar articles

15. Midtrimester preterm prelabour rupture of membranes (PPROM): expectant management or amnioinfusion for improving perinatal outcomes (PPROMEXIL - III trial).
van Teeffelen AS, van der Ham DP, Willekes C, Al Nasiry S, Nijhuis JG, van Kuijk S, Schuyt E, Mulder TL, Franssen MT, Oepkes D, Jansen FA, Woiski MD, Bekker MN, Bax CJ, Porath MM, de Laat MW, Mol BW, Pajkrt E.
BMC Pregnancy Childbirth. 2014 Apr 4;14:128. doi: 10.1186/1471-2393-14-128.
PMID: 24708702 Free PMC Article
Similar articles

16. Risk factors for uterine atony/postpartum hemorrhage requiring treatment after vaginal delivery.
Wetta LA, Szychowski JM, Seals S, Mancuso MS, Biggio JR, Tita AT.
Am J Obstet Gynecol. 2013 Jul;209(1):51.e1-6. doi: 10.1016/j.ajog.2013.03.011. Epub 2013 Mar 15.
PMID: 23507549 Free PMC Article
Similar articles

17. Induction of labor versus expectant management in women with preterm prelabor rupture of membranes between 34 and 37 weeks: a randomized controlled trial.
van der Ham DP, Vijgen SM, Nijhuis JG, van Beek JJ, Opmeer BC, Mulder AL, Moonen R, Groenewout M, van Pampus MG, Mantel GD, Bloemenkamp KW, van Wijngaarden WJ, Sikkema M, Haak MC, Pernet PJ, Porath M, Molkenboer JF, Kuppens S, Kwee A, Kars ME, Woiski M, Weinans MJ, Wildschut HI, Akerboom BM, Mol BW, Willekes C; PPROMEXIL trial group.
PLoS Med. 2012;9(4):e1001208. doi: 10.1371/journal.pmed.1001208. Epub 2012 Apr 24.
PMID: 22545024 Free PMC Article
Similar articles

18. Combined spinal and epidural anaesthesia and maternal intrapartum temperature during vaginal delivery: a randomized **clinical trial**.
de Orange FA, Passini R Jr, Amorim MM, Almeida T, Barros A.

RU 23:11 09.01.2019

pubmed - ничего не на | Induction of labour | Google Переводчик | преждевременный ра | Преждевременный ра | ДМ Акушер-Гинеколог | ДЗ ДМ на 10.01.19г..pp

https://www.ncbi.nlm.nih.gov/pubmed/17617892

NCBI Resources How To Sign in to NCBI

PubMed.gov US National Library of Medicine National Institutes of Health

PubMed | prenatal rupture of membranes | Search

Advanced Help

Format: Abstract Send to

BMC Pregnancy Childbirth. 2007 Jul 6;7:11.

Induction of labour versus expectant management in women with preterm prelabour rupture of membranes between 34 and 37 weeks (the PPROMEXIL-trial).

van der Ham DP[1], Nijhuis JG, Mol BW, van Beek JJ, Opmeer BC, Bijlenga D, Groenewout M, Arabin B, Bloemenkamp KW, van Wijngaarden WJ, Wouters MG, Pernet PJ, Porath MM, Molkenboer JF, Derks JB, Kars MM, Scheepers HC, Weinans MJ, Woiski MD, Wildschut HI, Willekes C.

Author information

Abstract

BACKGROUND: Preterm prelabour rupture of the membranes (PPROM) is an important clinical problem and a dilemma for the gynaecologist. On the one hand, awaiting spontaneous labour increases the probability of infectious disease for both mother and child, whereas on the other hand induction of labour leads to preterm birth with an increase in neonatal morbidity (e.g., respiratory distress syndrome (RDS)) and a possible rise in the number of instrumental deliveries.

METHODS/DESIGN: We aim to determine the effectiveness and cost-effectiveness of immediate delivery after PPROM in near term gestation compared to expectant management. Pregnant women with preterm prelabour rupture of the membranes at a gestational age from 34+0 weeks until 37+0 weeks will be included in a multicentre prospective randomised controlled trial. We will compare early delivery with expectant monitoring. The primary outcome of this study is neonatal sepsis. Secondary outcome measures are maternal morbidity (chorioamnionitis, puerperal sepsis) and neonatal disease, instrumental delivery rate, maternal quality of life, maternal preferences and costs. We anticipate that a reduction of neonatal infection from 7.5% to 2.5% after induction will outweigh an increase in RDS and additional costs due to admission of the child due to prematurity. Under these assumptions, we aim to randomly allocate 520 women to two groups of 260 women each. Analysis will be by intention to treat. Additionally a cost-effectiveness analysis will be performed to evaluate if the cost related to early delivery will outweigh those of expectant management. Long term outcomes will be evaluated using modelling.

DISCUSSION: This trial will provide evidence as to whether induction of labour after preterm prelabour rupture of membranes is an effective and cost-effective strategy to reduce the risk of neonatal sepsis. CONTROLLED CLINICAL TRIAL REGISTER: ISRCTN29313500.

PMID: 17617892 PMCID: PMC1934382 DOI: 10.1186/1471-2393-7-11

Full text links

Read free full text at BMC | PMC FREE Full text

Save items

Add to Favorites

Similar articles

Economic analysis comparing induction of labor and expectan [Acta Obstet Gynecol Scand. 2014]

An economic evaluation of planned immediate versus delayed birth for preterm pre [BJOG. 2017]

Immediate delivery compared with expectant management after preterm pre-lab [Lancet. 2016]

Review Planned early birth versus expectant managemer [Cochrane Database Syst Rev. 2006]

Review Antimicrobial therapy in expectant management of preterm premature [Lancet. 1995]

See reviews...

See all...

Cited by 5 PubMed Central articles

RU 23:07 09.01.2019

https://www.ncbi.nlm.nih.gov/pubmed/9465815

pubmed - ничего не найдено. | prenatal rupture of membran | Диалоги

https://www.ncbi.nlm.nih.gov/pubmed

Show additional filters

Search results

Items: 5

Filters activated: Clinical Trial, Humans. Clear all to show 138 items.

1. Single versus weekly courses of antenatal corticosteroids in preterm premature **rupture** of **membranes**.
   Lee MJ, Davies J, Guinn D, Sullivan L, Atkinson MW, McGregor S, Parilla BV, Hanlon-Lundberg K, Simpson L, Stone J, Wing D, Ogasawara K, Muraskas J.
   Obstet Gynecol. 2004 Feb;103(2):274-81.
   PMID: 14754695
   Similar articles

2. Premature **rupture** of **membranes** at 34 to 37 weeks' gestation: aggressive versus conservative management.
   Naef RW 3rd, Allbert JR, Ross EL, Weber BM, Martin RW, Morrison JC.
   Am J Obstet Gynecol. 1998 Jan;178(1 Pt 1):126-30.
   PMID: 9465815
   Similar articles

3. Randomized trial of endovaginal ultrasound in preterm premature **rupture** of **membranes**.
   Carlan SJ, Richmond LB, O'Brien WF.
   Obstet Gynecol. 1997 Mar;89(3):458-61.
   PMID: 9052605
   Similar articles

4. Group B Streptococcus and preterm premature **rupture** of **membranes**: a randomized, double-blind **clinical trial** of antepartum ampicillin.
   Grable IA, Garcia PM, Perry D, Socol ML.
   Am J Obstet Gynecol. 1996 Oct;175(4 Pt 1):1036-42.
   PMID: 8885772
   Similar articles

5. [Quantitative determination of fetal fibronectin in cervical smears: a new marker for evaluating the risk of premature labor].

Terms] OR "membranes"[All Fields])
AND ("chorioamnionitis"[MeSH Terms]

Search

See more...

Recent Activity

Turn Off Clear

- Premature rupture of membranes at 34 to 37 weeks' gestation: aggressive versus c PubMed
- prenatal rupture of membranes chorioamnionitis AND (Clinical Tria... (PubMed
- prenatal rupture of membranes chorioamnionitis AND (Clinical Tria... (PubMed
- prenatal rupture of membranes chorioamnionitis AND (Clinical Tria... (PubMed
- prenatal rupture of membranes chorioamnionitis (138) PubMed

See more...

RU 23:22 09.01.2019