

PARACETAMOL

Student: Chyndyn-ool Buian-Badyrgy
Solangyevich

Group: F-22(1)



Indications of use

- Pain of mild to moderate intensity
different genesis
- Fever in infectious and inflammatory
diseases
- headache
- Neuralgia



Contraindications

- Chronic alcoholism
- Hypersensitivity to paracetamol

Dosage

Inside or rectally in adults and adolescents weighing more than 60 kg is used in a single dose of 500 mg, the multiplicity of reception - up to 4 times / day. The maximum duration of treatment - 5-7 days.

The maximum doses: single - 1 g daily - 4 g

Single doses for intake for children aged 6-12 years - 250-500 mg, 1-5 years - 120-250 mg, from 3 months to 1 year - 60-120 mg, up to 3 months - 10 mg / kg.

Single doses for rectal use in children aged 6-12 years - 250-500 mg, 1-5 years - 125-250 mg.

An anatomical illustration of a human torso, showing the ribcage, spine, and internal organs. The liver is highlighted in a bright pink color, positioned in the upper right quadrant of the abdomen. The rest of the body is rendered in a semi-transparent, light blue/purple hue.

Overdose


liver failure

Side effects

An anatomical illustration of the human digestive system, including the liver, stomach, and intestines, overlaid on a blue-tinted human torso. The liver is shown in the upper right quadrant, the stomach is in the upper left, and the coiled small and large intestines occupy the lower abdomen.

- ❖ rarely – dyspepsia
- ❖ long-term use at high doses - hepatotoxic effects

- 
- A microscopic view of red blood cells (erythrocytes) in a blood vessel. The cells are biconcave discs, appearing as reddish-orange spheres with a darker center. They are scattered throughout the field, with some in sharp focus and others blurred in the background. The overall color is a deep red, and the lighting creates a sense of depth and movement.
- Thrombocytopenia
 - leukopenia
 - pancytopenia
 - neutropenia
 - agranulocytosis

- 
- A close-up photograph of human skin showing a significant allergic reaction. The skin is covered in a dense, irregular pattern of bright red, raised, and inflamed patches, characteristic of urticaria (hives). The background is a blurred, warm-toned skin surface.
- skin rash
 - Itch
 - urticaria

Terms and conditions of storage

A pharmacist with blonde hair in a braid, wearing glasses and a white lab coat, is standing in a pharmacy. She is holding a tablet in her left hand and pointing with her right hand towards the shelves of medicine. The shelves are filled with various colorful boxes and bottles of pharmaceuticals.

Store in tightly closed containers in a dry, dark place at room temperature.

Shelf life of paracetamol Tablets - 3 years;
slurry - 2 years.

Thank you for attention 😊

