

# PARACETAMOL

Student: Chyndyn-ool Buian-Badyrgy  
Solangyevich  
Group: F-22(1)

# Indications of use



- Pain of mild to moderate intensity  
different genesis
- Fever in infectious and inflammatory  
diseases
- headache
- Neuralgia





# Contraindications

- Chronic alcoholism
- Hypersensitivity to paracetamol

# Dosage

Inside or rectally in adults and adolescents weighing more than 60 kg is used in a single dose of 500 mg, the multiplicity of reception - up to 4 times / day. The maximum duration of treatment - 5-7 days.

The maximum doses: single - 1 g daily - 4 g

Single doses for intake for children aged 6-12 years - 250-500 mg, 1-5 years - 120-250 mg, from 3 months to 1 year - 60-120 mg, up to 3 months - 10 mg / kg.

Single doses for rectal use in children aged 6-12 years - 250-500 mg, 1-5 years - 125-250 mg.



An anatomical illustration of a human torso, showing the internal organs. The liver is highlighted in a bright pink color, contrasting with the blue and purple tones of the rest of the body. The liver is located in the upper right quadrant of the abdomen. The text "Overdose" is written in white, bold, sans-serif font across the upper part of the image, and "liver failure" is written in the same style across the lower part. The background is a dark purple gradient.

# Overdose


# liver failure




# Side effects



- ❖ rarely – dyspepsia
- ❖ long-term use at high doses - hepatotoxic effects

- 
- A microscopic view of a blood smear showing numerous red blood cells. The cells are biconcave discs, appearing as reddish-orange circles with darker centers. They are distributed across the field of view, with some in sharp focus and others blurred in the background. The overall color is a deep red with some lighter, yellowish-orange areas where the cells are more concentrated or where the background is visible.
- Thrombocytopenia
  - leukopenia
  - pancytopenia
  - neutropenia
  - agranulocytosis



- 
- The background of the slide is a close-up photograph of human skin affected by a rash. The skin is covered with numerous red, raised, and inflamed lesions, which are characteristic of urticaria (hives). The lesions vary in size and are distributed across the visible skin area.
- skin rash
  - Itch
  - urticaria



# Terms and conditions of storage



Store in tightly closed containers in a dry, dark place at room temperature.

Shelf life of paracetamol Tablets - 3 years;  
slurry - 2 years.

Thank you for attention 🥰

