



Skin and soft tissue infections

Shingles, Molluscum Contagiosum, Staphylococcal skin syndrome, scabies, lice infection, erythrasma

Повестка


Раздел 1

Раздел 2

Раздел 3

Раздел 4

Раздел 5





Введение

В PowerPoint можно создавать презентации и делиться своими материалами с другими, где бы они ни находились. Введите здесь нужный текст, чтобы начать работу. В этом шаблоне вы также можете добавлять рисунки, изображения и видеоролики. Сохраняйте презентации в OneDrive и открывайте их с компьютера, планшета или телефона.


Scabies

Mites *Sarcoptes scabiei*



Scabies


- Transmission
crowding, skin to skin contact and fomites
- The female mite burrows just beneath the skin
- in order to lay her eggs. She then dies.
- The eggs hatch into tiny mites that spread out over the skin and live for only about 30 days.



Scabies

CLASSIC SCABIES

- Small erythematous papules
- “Knots on a rope”
- Pruritic
- Severe and worse at night
- Fingers web spaces, flexor surfaces of wrist, elbow, axillary folds, beltline, lower buttocks, genitalia




Scabies

CRUSTED SCABIES

- Norwegian scabies
- Crusting, scaling fissuring affecting an older, immunosuppressed adult
- Higher mite burden
- Transmission via fomites
- Hands, feet, scalp



Scabies



- Diagnosis
- Clx – history and appearance of the rash
- Microscopy
- Treatment
- Permethrim
- Lindane
- Ivermectim

Head Lice
Body Lice
Pubic Lice

Pediculosis

Pediculosis humanus capitis

Pediculosis humanus corporis

Pediculosis pubis

Pediculosis ciliaris



Head louse

Photo: Lorenza Beati
Used with permission



Body louse

Photo: James Gathany
Public Health Image Library (PHIL)
Public domain



Crab louse

Photo: Matt Bertone
Used with permission

Compiled by: Slavash Taravati, UC IPM program
Compiled by: Slavash Taravati, UC IPM program



Head Lice

- Children, females, European
- Direct contact or fomites
- Nits firmly “cemented” to human hair
- White spots of nits can be mistaken for dandruff
- Unlike dandruff, the nits cannot be brushed off





Body Lice


- Poverty, poor hygiene, crowding
- Direct contact and clothing
- Lays eggs in seams of clothing
- Can live up to 3 days without feeding on host





Pubic Lice

- Sexual active, young adults and adolescents
- Sexual transmitted and fomites
- Contact with eyes can lead to Pediculosis ciliaris
- Generally smaller in size than the other types



Lice

Symptoms

- Itchy
- Excoriation
- Hyperpigmentation
- Lymphadenopathy

Bacteria transmitted by the body louse

- *Rickettsia prowazekii*
- *Borrelia recurrents*
- *Borrelia quintana*

Lice

Diagnosis

- Head lice or nits are usually on the scalp and nape of the neck and over the ears. Adult lice are approximately the same size as a sesame seed.
- Body lice are more difficult to find, but they usually can be detected in the seams of underwear.
- Pubic lice are found on the skin and hair of the pubic area or on the eyelashes

Treatment

- Inspection and remove
- Permethim
- Refractory treatment: oral Ivermectin




Herpes zoster Shingles

Varicella zoster virus



Shingles

- Due to reactivation of the VZV from dorsal root ganglion nerves
- *the condition is preceded by several days of radicular pain with hyperaesthesia*
- *unilateral patchy rash in one or two contiguous dermatomes*
- *intense erythema with papules in affected skin*
- *later crusting and separation of scabs after 10–14 days, often with depigmentation*
- *regional lymphadenopathy*



Shingles

Herpes Zoster oticus the trigeminal nerve

- Ramsay Hunt Syndrome
 - Ipsilateral facial paralysis
 - Ear pain
 - Vesicles in the auditory canal and auricle

Herpes Zoster ophthalmicus the facial nerve

- Conjunctivitis, scleritis, episcleritis, keratitis, glaucoma, retinitis
- Argyll-Robertson pupils



Shingles

Diagnostic

- RCR for detection of viral DNA
- Direct fluorescent antibody
- Tzanck smear

Treatment

- <72 hrs – valacyclovir, acyclovir
- Post-herpetic neuralgia
 - Amitriptyline, pregabalin, gabapentin

Molluscum contagiosum

Molluscum contagiosum virus

Poxvirus



Molluscum contagiosum

- Children and sexually active adults
- Painless
- Incubation period
 - between 2-6 weeks
- Persist for months
- In any part of the body except palms and soles
- Sometimes pruritic

Transmission

- Direct skin-to-skin contact
- Autoinoculation
 - “kissing lesions”
- Fomites
 - sharing towels and bath toys
- Through water
 - Swimming pool

Molluscum contagiosum


- Dome-shaped papules with umbilication
- 2-3mm in diameter
- Pink-white to flesh colored
- Single or multiple (more common)
- Hemispherical up to 5 mm



Molluscum contagiosum

Diagnosis

- Clinical
- Histology
 - Henderson – Peterson bodies



Treatment

- Self-limiting
- Cryotherapy, Cantharidin, Curretage, Imiquimod, Topical retinoids

Staphylococcal scalded skin syndrome

Staphylococcus aureus



SSSS

- is a bacterial toxin-mediated skin disorder that primarily affects young children
- generally from bullous Impetigo
- occurs when exotoxins produced by *Staphylococcus aureus* undergo hematogenous dissemination to the skin
- diffuse skin pain and erythema as well as superficial blistering and desquamation
- fever, irritability, and poor oral intake.
- The desquamation phase lasts 2 to 4 days and is followed by complete healing, without scarring



Diagnosis

- Clx
- Skin examination
 - Flaccid bullae, superficial desquamation, and shallow erosions
 - Absent mucous membrane involvement
 - Evidence of concurrent cutaneous, conjunctival, or internal staphylococcal infection
 - Positive Nikolsky sign

Treatment

- Intravenous antimicrobials
 - oxacillin or nafcillin.
- Isolation in an incubator
- Nontraumatic skin care
 - emollients (sterile petrolatum, paraffin oil);
- the shedding epidermis must be conserved as a “biologic dressing”


Erythrasma

Corynebacterium minutissimum



Erythrasma

- chronic superficial skin infection
- Superficial reddish-brown scaly patches
- Enlarges peripherally
- Mild infection but tends to chronicity if untreated
- Coral pink fluorescence with Wood's light
- Common sites: groin (especially men), axillae, submammary, toe webs





Erythrasma

Diagnosis

- Appearance
- Fluoresces coral red with Wood's lamp
- KOH exam for dermatophyte (that can co-exist)
- Gram stain: g+ filaments and rods



Erythrasma

Treatment

- Topical imidazole e.g. miconazole or erythromycin 2% gel
- Oral roxithromycin or erythromycin
- Loose fitting clothing and antibacterial wash may prevent recurrence

Tinea versicolor



